arXiv:1504.06452v2 [math-ph] 17 Jun 2015

Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{\mathcal{M}}_{g,n}$

Marco Bertola, Boris Dubrovin, Di Yang

June 18, 2015

Abstract

We derive an explicit generating function of correlations functions of an arbitrary tau-function of the KdV hierarchy. In particular applications, our formulation gives closed formulæ of a new type for the generating series of intersection numbers of ψ-classes as well as of mixed ψ- and κ-classes in full genera.

Keywords. KdV hierarchy; tau-function; wave function; correlation function; intersection number.

Contents

1 Introduction and results **2**

2 A brief reminder of the KdV hierarchy **10**

3 One-point correlation functions of the KdV hierarchy **12**

4 Multi-point correlation functions of the KdV hierarchy **14**

4.1 Generating function of two-point correlation functions 15

4.2 Generating functions of multi-point correlation functions 16

5 On string equation and Kac–Schwarz operator **18**

6 Application to the Witten–Kontsevich tau-function **19**

7 Application to higher Weil–Petersson volumes **23**

7.1 Proof of Theorem 1.14 24

7.2 Examples of computations of higher Weil–Petersson volumes 27

8 Further remarks **32**

A Some useful formulæ **34**

B Generalized Kac–Schwarz operator and higher Weil–Petersson volumes **34**

C Tables of some intersection numbers **38**

1 Introduction and results

The famed Korteweg–de Vries (KdV) equation

$$u_t = uu_x + \frac{1}{12}u_{xxx}, \tag{1.1}$$

has long been known to be *integrable*. It belongs to an infinite family of pairwise commuting nonlinear evolutionary PDEs called *the KdV hierarchy*. The hierarchy can be described in terms of isospectral deformations of the Lax operator

$$L = \partial_x^2 + 2\,u(x). \tag{1.2}$$

Namely, the k-th equation of the KdV hierarchy, in the normalization of the present paper, reads

$$L_{t_k} = [A_k, L], \tag{1.3}$$

$$A_k := \frac{1}{(2k+1)!!}\left(L^{\frac{2k+1}{2}}\right)_+, \quad k \geq 0. \tag{1.4}$$

Here, the independent variables $t_0, t_1, t_2, ...$ are called *times*. The symbol $\left(L^{\frac{2k+1}{2}}\right)_+$ stands for the differential part of the pseudo-differential operator $L^{\frac{2k+1}{2}}$, see e.g. the book [16] for details. The $k = 1$ equation of (1.3) coincides with (1.1). As customary in the literature, we shall identify t_0 with the spatial variable x.

The notion of *tau-function* for the KdV hierarchy was introduced by the Kyoto school [50, 27, 15] during the 1970s–1980s. In 1991, E. Witten, in his study of two-dimensional quantum gravity [52], conjectured that the generating function of the intersection numbers of ψ-classes on the Deligne–Mumford moduli spaces $\overline{\mathcal{M}}_{g,n}$ of stable algebraic curves is a tau-function of the KdV hierarchy. Witten's conjecture was later proved by M. Kontsevich [34]; see [33, 49, 44] for several alternative proofs. Moreover the so-called "tau structures" of KdV-like hierarchies became one of the central subjects in the study of the deep relation between integrable hierarchies and Gromov–Witten invariants [22, 20, 21].

The original motivations of the Witten's conjecture identify the tau-function $\tau = \tau(t_0, t_1, t_2, \ldots)$ of a particular solution to the KdV hierarchy with the partition function of 2D quantum gravity. The time variables t_0, t_1, t_2, $\ldots$ are identified with coupling constants associated with observables $\tau_0 = 1$, τ_1, τ_2, $\ldots$ of the quantum theory. Thus, the quantum correlators of 2D quantum gravity are just logarithmic derivatives of the tau-function

$$\langle \tau_{k_1}\tau_{k_2}\ldots\tau_{k_n}\rangle = \left.\frac{\partial^n \log\tau(t_0,t_1,t_2,\ldots)}{\partial t_{k_1}\partial t_{k_2}\ldots\partial t_{k_n}}\right|_{t_0=t_1=t_2=\ldots=0}. \tag{1.5}$$

Tau-functions of some other solutions to the KdV hierarchy along with the corresponding correlators of the form (1.5) proved to be of interest for other applications, in particular to the study of topology of Deligne–Mumford moduli spaces (see below). The main goal of this paper is to provide a simple algorithm, in the framework of the theory of Lax operator and its eigenfunctions, for computation of n–point correlators of an arbitrary solution to the KdV hierarchy.

Let us begin with basics of the theory of KdV tau-functions in the version of M. Sato *et al.* Denote by $\mathbf{t} = (t_0, t_1, t_2, \dots)$ the infinite vector of time variables. Let $\mathcal{B} = \mathbb{C}[[t_k,\, k = 0, 1, 2, \dots]]$ be the Bosonic Fock space. A *Sato tau-function* $\tau(\mathbf{t})$ of the KdV hierarchy is an element in $\mathcal{B}$ satisfying the Hirota bilinear identities

$$\operatorname*{res}_{z=\infty} \tau(\mathbf{t} - [z^{-1}])\, \tau(\widetilde{\mathbf{t}} + [z^{-1}]) \exp\left(\sum_{j\geq 1} \frac{t_j - \widetilde{t}_j}{(2j+1)!!} z^{2j+1} \right) z^{2p}\, dz = 0, \qquad \forall\, \mathbf{t},\, \widetilde{\mathbf{t}},\, p = 0, 1, 2, \dots. \tag{1.6}$$

Here $\mathbf{t} - [z^{-1}] := \left(t_0 - z^{-1}, \dots, t_k - \frac{(2k-1)!!}{z^{2k+1}}, \dots \right)$. The residue is understood in a formal way, namely, as (minus) the coefficient of z^{-1} in the formal expansion at $z = \infty$. Given an arbitrary tau-function $\tau(\mathbf{t})$, then $u(\mathbf{t}) = \partial_x^2 \log \tau(\mathbf{t})$ is a solution of the KdV hierarchy (1.3).

Conversely, let $u(\mathbf{t})$ be an arbitrary (formal) solution of the KdV hierarchy (1.3); then there exists [16] a tau-function $\tau(\mathbf{t})$ such that $\partial_x^2 \log \tau(\mathbf{t}) = u(\mathbf{t})$. The tau-function of $u(\mathbf{t})$ is uniquely determined up to a *gauge freedom*

$$\tau(\mathbf{t}) \mapsto \exp\Big(\alpha_{-1} + \sum_{j\geq 0} \alpha_j\, t_j \Big) \tau(\mathbf{t}) \tag{1.7}$$

where the coefficients α_j, $j \geq -1$ are arbitrary constants.

Let $\tau(\mathbf{t})$ be any tau-function of $u(\mathbf{t})$. Define the *wave* and *dual wave functions* by

$$\psi(z; \mathbf{t}) = \frac{\tau(\mathbf{t} - [z^{-1}])}{\tau(\mathbf{t})} \mathrm{e}^{\vartheta(z;\mathbf{t})}, \quad \psi^*(z; \mathbf{t}) = \psi(-z; \mathbf{t}) = \frac{\tau(\mathbf{t} + [z^{-1}])}{\tau(\mathbf{t})} \mathrm{e}^{-\vartheta(z;\mathbf{t})}, \tag{1.8}$$

where the phase ϑ is given by

$$\vartheta(z; \mathbf{t}) := \sum_{j=0}^{\infty} t_j \frac{z^{2j+1}}{(2j+1)!!}. \tag{1.9}$$

The gauge freedom (1.7) affects $\psi(z; \mathbf{t})$ by a multiplicative factor of the form

$$g(z) = \exp\left(-\sum_{k=0}^{\infty} \frac{\alpha_k \cdot (2k-1)!!}{z^{2k+1}} \right) = 1 + \mathcal{O}(z^{-1}), \qquad z \to \infty. \tag{1.10}$$

The wave functions are (formal) eigenfunctions of the Lax operator

$$L\, \psi = z^2\, \psi, \quad L\, \psi^* = z^2\, \psi^*. \tag{1.11}$$

Their dependence on time variables is specified by the following compatible system

$$\psi_{t_k} = A_k \psi, \quad \psi^*_{t_k} = -A_k \psi^*, \quad k \geq 0. \tag{1.12}$$

where the differential operators A_k are defined in (1.4). As $z \to \infty$ their (formal) asymptotic behaviours are

$$\psi(z; \mathbf{t}) = \big(1 + \mathcal{O}(z^{-1})\big)\, \mathrm{e}^{\vartheta(z;\mathbf{t})}, \quad \psi^*(z; \mathbf{t}) = \big(1 + \mathcal{O}(z^{-1})\big)\, \mathrm{e}^{-\vartheta(z;\mathbf{t})}. \tag{1.13}$$

Also they satisfy an infinite system of bilinear relations

$$\operatorname*{res}_{z=\infty} \psi(z; \mathbf{t})\, \psi^*(z; \widetilde{\mathbf{t}})\, z^{2p}\, dz = 0, \qquad \forall\, \mathbf{t},\, \widetilde{\mathbf{t}},\, p = 0, 1, \dots. \tag{1.14}$$

See e.g. [16], Thm. 6.3.8 for proofs of these statements. Depending on the context, the wave and dual wave functions (as well as the τ-function) can be defined analytically following the approach of G. Segal and G. Wilson [51]; also see our Remark 4.2 below.

Definition 1.1. *For any tau-function $\tau(\mathbf{t})$ of the KdV hierarchy, we call the functions*

$$\langle\langle \tau_{k_1}\tau_{k_2}\dots\tau_{k_n}\rangle\rangle(\mathbf{t}) := \frac{\partial^n \log\tau}{\partial t_{k_1}\dots\partial t_{k_n}}(\mathbf{t}), \quad n\geq 1 \tag{1.15}$$

and the numbers

$$\langle \tau_{k_1}\tau_{k_2}\dots\tau_{k_n}\rangle := \langle\langle \tau_{k_1}\tau_{k_2}\dots\tau_{k_n}\rangle\rangle(\mathbf{t}=0) \tag{1.16}$$

*the n-**point correlation functions** and the n-**point correlators** of the tau-function respectively. For every $n\geq 1$ we define the generating function of n-point correlation functions by*

$$F_n(z_1,\dots,z_n;\mathbf{t}) := \sum_{k_1,\dots,k_n=0}^{\infty} \langle\langle \tau_{k_1}\dots\tau_{k_n}\rangle\rangle(\mathbf{t})\,\frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}}. \tag{1.17}$$

Evaluating it at $\mathbf{t}=0$

$$F_n(z_1,\dots,z_n) := F_n(z_1,\dots,z_n;\mathbf{0}) \tag{1.18}$$

one obtains generating functions of n-point correlators of $\tau(\mathbf{t})$.

The notation (1.15) is borrowed from the literature in quantum gravity [52].

We are now in a position to formulate the first main result of the paper.

Theorem 1.2. *Let τ be any tau-function of the KdV hierarchy and let ψ,ψ^* be defined by (1.8). The generating function of one-point correlation functions has the following expression*

$$F_1(z;\mathbf{t}) = \frac{1}{2}\mathrm{Tr}\left(\Psi^{-1}(z)\Psi_z(z)\sigma_3\right) - \vartheta_z(z) \tag{1.19}$$

where

$$\Psi(z;\mathbf{t}) = \begin{pmatrix} \psi(z;\mathbf{t}) & \psi^*(z;\mathbf{t}) \\ -\psi_x(z;\mathbf{t}) & -\psi_x^*(z;\mathbf{t}) \end{pmatrix} \tag{1.20}$$

and $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ is the Pauli matrix.

The proof is in Sect. 3. The identity (1.19) gives a unique reconstruction, up to a multiplicative constant, of tau-function from the knowledge of the corresponding wave functions and thus provides a converse to Sato's result.

Remark 1.3. *The reader conversant with the theory of isomonodromic deformation will observe similarity with the Jimbo–Miwa–Ueno definition of isomonodromic tau-function [29, 7].*

Remark 1.4. *The theory developed by Sato et al. gives a formal expansion of τ by using the infinite Grassmannian approach (see for example in [28, 5, 55] for details)*

$$\tau = \sum_{\mu\in\mathbb{Y}} \pi_\mu\, \mathfrak{s}_\mu \tag{1.21}$$

which provides derivatives of τ at $\mathbf{t}=0$ in terms of Plücker coordinates. Here, $\mathbb{Y}$ denotes the set of Young diagrams, π_μ are Plücker coordinates of a point in the Sato Grassmannian, $\mathfrak{s}_\mu=\mathfrak{s}_\mu(T)$ are Schur polynomials in $T=(T_1,T_2,\dots)$ with $t_k=-(2k+1)!!\,T_{2k+1}$. However, the representation (1.21) does not help much with computation of logarithmic derivatives of tau-function.

A well-known formula (see e.g. [3], pag. 389), expressing the generating function of $\langle\langle\tau_0\tau_j\rangle\rangle$, can be easily obtained as an immediate corollary of Theorem 1.2:

Corollary 1.5. *The following equality holds true:*

$$1+\sum_{j=0}^{\infty}\frac{(2j+1)!!}{z^{2j+2}}\langle\langle\tau_0\tau_j\rangle\rangle=\psi(z)\,\psi^*(z). \tag{1.22}$$

It has been shown in [18, 3] that any pair of wave and dual wave functions ψ,ψ^* satisfy

$$\psi(z)\,\psi^*(z)=1+\sum_{j=0}^{\infty}\frac{Res_\partial L^{\frac{2j+1}{2}}}{z^{2j+2}}. \tag{1.23}$$

Hence from (1.23) and (1.22) we have $\langle\langle\tau_0\tau_j\rangle\rangle=\frac{1}{(2j+1)!!}Res_\partial L^{\frac{2j+1}{2}}$, $j\geq 0$.

Our next result expresses the generating function of two-point correlation functions in terms of wave and dual wave functions.

Proposition 1.6. *Let τ be a tau-function of the KdV hierarchy and let ψ,ψ^* defined by (1.8). The generating function of two-point correlation functions has the following expression*

$$F_2(z,w;\mathbf{t})=\frac{\frac{1}{2}\mathcal{R}_x(w)\mathcal{R}_x(z)-\mathcal{R}(w)\mathcal{R}(z)\chi(z)\chi(-z)-\mathcal{R}(z)\mathcal{R}(w)\chi(w)\chi(-w)-(z^2+w^2)}{(z^2-w^2)^2} \tag{1.24}$$

where $\mathcal{R}=\mathcal{R}(z;\mathbf{t})=\psi(z;\mathbf{t})\,\psi^*(z;\mathbf{t})$, $\chi=\chi(z;\mathbf{t})=\partial_x\log\psi(z;\mathbf{t})$; the explicit dependence on $\mathbf{t}$ of the functions $\mathcal{R}(z;\mathbf{t})$, $\mathcal{R}(w;\mathbf{t})$ etc. is omitted from (1.24).

The proof is in Section 4.1. Note that both $\mathcal{R}$ and χ are intended as formal Laurent series in z^{-1} whose coefficients are differential polynomials in u. The formula (1.24) was also derived in [22] through the Hadamard–Seeley expansion of $\mathrm{e}^{\varepsilon L}$, $\varepsilon\to 0$.

Now we formulate the result for n-point correlation functions ($n\geq 2$); it includes the case $n=2$ but we have preferred to highlight it separately in the above Prop. 1.6 for clarity's sake.

Theorem 1.7 (Main Theorem)**.** *Let τ be any tau-function of the KdV hierarchy and let $\psi=\psi(z;\mathbf{t}),\psi^*=\psi^*(z;\mathbf{t})$ be defined by (1.8). Let $\mathcal{R}=\mathcal{R}(z;\mathbf{t})$ be as above. Denote Θ the matrix given by one of the following equivalent expressions*

$$\Theta(z;\mathbf{t}) = \frac{1}{2}\begin{bmatrix} -\mathcal{R}_x & -2\mathcal{R} \\ \mathcal{R}_{xx}-2(z^2-2\,u)\mathcal{R} & \mathcal{R}_x \end{bmatrix} \tag{1.25}$$

$$= \frac{1}{2}\begin{bmatrix} -(\psi\,\psi^*)_x & -2\,\psi\,\psi^* \\ 2\,\psi_x\,\psi^*_x & (\psi\,\psi^*)_x \end{bmatrix} \tag{1.26}$$

$$= z\,\Psi(z;\mathbf{t})\,\sigma_3\,\Psi^{-1}(z;\mathbf{t}) \tag{1.27}$$

where the matrix $\Psi(z;\mathbf{t})$ is defined in (1.20). Then the generating function of n-point correlation functions with $n \geq 2$ has the following expression

$$F_n(z_1,\dots,z_n;\mathbf{t}) = -\frac{1}{n}\sum_{r\in S_n}\frac{\operatorname{Tr}\left(\Theta(z_{r_1})\cdots\Theta(z_{r_n})\right)}{\prod_{j=1}^n (z_{r_j}^2 - z_{r_{j+1}}^2)} - \delta_{n,2}\frac{z_1^2+z_2^2}{(z_1^2-z_2^2)^2}. \tag{1.28}$$

Here $\delta_{n,2}$ is the Kronecker delta, S_n denotes the group of permutations of $\{1,\dots,n\}$, and for a permutation $r=[r_1,\dots,r_n]$, r_{n+1} is understood as r_1.

The proof is in Sect. 4.

Remark 1.8. *The reader can verify that the right side of (1.28) is regular on the diagonals $z_i = z_j$.*

Example 1.9. *The next example is $n=3$ for which the theorem above yields (on account of the cyclicity of the trace)*

$$F_3(z_1,z_2,z_3;\mathbf{t}) = -\frac{\operatorname{Tr}\left(\Theta(z_1)\Theta(z_2)\Theta(z_3)\right) - \operatorname{Tr}\left(\Theta(z_2)\Theta(z_1)\Theta(z_3)\right)}{(z_1^2-z_2^2)(z_2^2-z_3^2)(z_3^2-z_1^2)}. \tag{1.29}$$

Remark 1.10. *We have been recently made aware of an interesting preprint from Si-Qi Liu [39], where the $n=2,3$ cases of Theorem 1.7 have also been obtained independently. Jian Zhou also privately communicated a different nice proof of Theorem 1.7 [56].*

Application to intersection numbers of ψ-classes on $\overline{\mathcal{M}}_{g,n}$. Recall that $\overline{\mathcal{M}}_{g,n}$ denotes the Deligne–Mumford moduli space of stable curves of genus g with n marked points; let us also recall

Witten's conjecture (Kontsevich's theorem): *The partition function of 2D quantum gravity*

$$Z(\mathbf{t}) = \exp\left(\sum_{n=0}^{\infty}\frac{1}{n!}\sum_{k_1,\dots,k_n\geq 0}\langle\tau_{k_1}\cdots\tau_{k_n}\rangle\, t_{k_1}\cdots t_{k_n}\right) \tag{1.30}$$

is a tau-function of the KdV hierarchy (1.3). Here $\langle\tau_{k_1}\dots\tau_{k_n}\rangle$ are intersection numbers of ψ-classes over the Deligne–Mumford moduli spaces

$$\langle\tau_{k_1}\dots\tau_{k_n}\rangle := \begin{cases} \displaystyle\int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{k_1}\dots\psi_n^{k_n}, & \text{if } g := \tfrac{k_1+\dots+k_n-n+3}{3} \text{ is a nonnegative integer,} \\ 0, & \text{otherwise.} \end{cases} \tag{1.31}$$

In the above integrals, ψ_i is the first Chern class of the i-th tautological line bundle over $\overline{\mathcal{M}}_{g,n}$.

The partition function $Z(\mathbf{t})$ is now generally known as the *Witten–Kontsevich tau-function*. Witten's conjecture implies that $u^{WK}(\mathbf{t}) := \partial_x^2 \log Z(\mathbf{t})$ is a particular solution of (1.3), for which reason we call it the *Witten–Kontsevich solution*, also known as *the topological solution*. It can be specified by the initial data

$$u^{WK}(\mathbf{t})|_{t_0=x,\ t_{\geq 1}=0} = x. \tag{1.32}$$

Applying Theorem 1.2 and Theorem 1.7 to the Witten–Kontsevich tau-function, we obtain

Theorem 1.11. *Let $M(z)$ denote the following matrix-valued formal series*

$$M(z) = \frac{1}{2}\begin{pmatrix} -\sum_{g=1}^{\infty} \frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!} z^{-6g+4} & -2\sum_{g=0}^{\infty} \frac{(6g-1)!!}{24^g\cdot g!} z^{-6g} \\ 2\sum_{g=0}^{\infty} \frac{6g+1}{6g-1}\frac{(6g-1)!!}{24^g\cdot g!} z^{-6g+2} & \sum_{g=1}^{\infty} \frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!} z^{-6g+4} \end{pmatrix}. \tag{1.33}$$

The generating functions of n-point intersection numbers (1.31)

$$F_n^{WK}(z_1,\dots,z_n) := \sum_{k_1,\dots,k_n=0}^{\infty} \langle \tau_{k_1}\dots\tau_{k_n}\rangle \frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}}, \quad n\geq 1 \tag{1.34}$$

are given by the following formulæ:

$$F_1^{WK}(z) = \sum_{g=1}^{\infty} \frac{(6g-3)!!}{24^g\cdot g!} z^{-(6g-2)}, \tag{1.35}$$

$$F_n^{WK}(z_1,\dots,z_n) = -\frac{1}{n}\sum_{r\in S_n} \frac{\mathrm{Tr}\,(M(z_{r_1})\cdots M(z_{r_n}))}{\prod_{j=1}^n (z_{r_j}^2 - z_{r_{j+1}}^2)} - \delta_{n,2}\frac{z_1^2+z_2^2}{(z_1^2-z_2^2)^2}, \quad n\geq 2. \tag{1.36}$$

The expression (1.35) is equivalent to the well-known formula (for example see in [25, 48]):

$$\langle \tau_{3g-2}\rangle = \frac{1}{24^g\cdot g!}, \quad g\geq 1. \tag{1.37}$$

The formula (1.36) seems to be new.

Remark 1.12. *In [48], A. Okounkov provided an expression for different generating functions of the same intersection numbers. The relationship between the two approaches is the following*

Our approach	*Okounkov's approach*
$\displaystyle\sum_{k_1,\dots,k_n=0}^{\infty} \frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}} \langle \tau_{k_1}\dots\tau_{k_n}\rangle$	$\displaystyle\sum_{k_1,\dots,k_n=0}^{\infty} x_1^{k_1}\cdots x_n^{k_n} \langle \tau_{k_1}\dots\tau_{k_n}\rangle$

(1.38)

It should be apparent that the two generating functions are related by a (formal) multiple Laplace transform. Okounkov's approach produced integral expressions (but reducing to explicit formulæ for $n\leq 3$). On the other hand the formula (1.36) is absolutely explicit and allows an efficient computation of the n-point numbers also for very high genera. See Section 8 below for more details.

Remark 1.13. *In [6] M. Bergere and B. Eynard introduced correlators associated to the Christoffel–Darboux kernel obtained from a general solution of a 2×2 linear ODE (see Definition 2.3 in [6]). Their construction was extended in [2] to Christoffel–Darboux kernels asociated with larger systems of ODEs. In the particular case of Airy kernel it can be shown that the Bergere–Eynard generating functions of the correlators, after a suitable normalization, reproduce intersection numbers of ψ-classes. Our generating function (1.36) obtained by restricting the general expression of the Main Theorem to the Witten–Kontsevich solution has a similar form to the generating function (2-9) of [6]. It is unclear to what kernels one should apply the Bergere–Eynard approach in order to derive, for example, the generating function of higher Weil–Petersson volumes (see below).*

Application to higher Weil–Petersson volumes. The *Main Theorem* also allows us to compute higher Weil–Petersson volumes, by which we mean integrals of mixed ψ- and κ-classes of the form

$$\langle \kappa_1^{d_1} \dots \kappa_\ell^{d_\ell} \tau_{k_1} \dots \tau_{k_n} \rangle_{g,n} := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{k_1} \dots \psi_n^{k_n} \kappa_1^{d_1} \dots \kappa_\ell^{d_\ell}. \tag{1.39}$$

Recall [1, 47] that κ-classes as elements of the Chow ring $A^*\left(\overline{\mathcal{M}}_{g,n}\right)$ are defined by

$$\kappa_i = f_*\left(\psi_{n+1}^{i+1}\right) \in A^i\left(\overline{\mathcal{M}}_{g,n}\right), \quad i \geq 0 \tag{1.40}$$

where $f : \overline{\mathcal{M}}_{g,n+1} \to \overline{\mathcal{M}}_{g,n}$ is the universal curve (the forgetful map). The class $\kappa_0 = 2g-2+n$ is just a constant, so it will not appear below. The integrals (1.39) take zero values unless

$$\sum_{j=1}^{n} k_j + \sum_{j=1}^{l} j\, d_j = 3g - 3 + n. \tag{1.41}$$

We denote by $Z^\kappa(\mathbf{t};\mathbf{s})$ the partition function of higher Weil–Petersson volumes

$$Z^\kappa(\mathbf{t};\mathbf{s}) = \exp\left(\sum_{g,n,\ell \geq 0} \frac{1}{n!} \sum_{d_1,\dots,d_\ell,k_1,\dots,k_n \geq 0} \langle \kappa_1^{d_1} \dots \kappa_\ell^{d_\ell} \tau_{k_1} \dots \tau_{k_n} \rangle_{g,n}\, t_{k_1} \cdots t_{k_n} \frac{s_1^{d_1} \dots s_\ell^{d_\ell}}{d_1! \cdots d_\ell!} \right), \tag{1.42}$$

where $\mathbf{s}$ denotes the infinite vector of independent variables $(s_1, s_2, \dots)$. It is a KdV tau-function of a family of solutions depending on the parameters $\mathbf{s}$ [32]. The $\mathbf{s} = \mathbf{0}$ evaluation of this function gives the Witten–Kontsevich tau-function: $Z(\mathbf{t}) = Z^\kappa(\mathbf{t};\mathbf{0})$. By $\psi^{WK}(z;\mathbf{t})$, $\psi^\kappa(z;\mathbf{t};\mathbf{s})$ denote the corresponding wave functions

$$\psi^{WK}(z;\mathbf{t}) := \frac{Z(\mathbf{t} - [z^{-1}])}{Z(\mathbf{t})} \mathrm{e}^{\vartheta(z;\mathbf{t})}, \quad \psi^\kappa(z;\mathbf{t};\mathbf{s}) := \frac{Z^\kappa(\mathbf{t} - [z^{-1}];\mathbf{s})}{Z^\kappa(\mathbf{t};\mathbf{s})} \mathrm{e}^{\vartheta(z;\mathbf{t})}. \tag{1.43}$$

Notations. (i) $\mathbb{Y}$ will denote the set of all partitions. (ii) For a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$, we denote by $\ell(\lambda) = card\{i = 1, 2, \dots \,|\, \lambda_i \neq 0\}$ the *length* of λ, by $|\lambda| = \sum_{i \geq 1} \lambda_i$ the *weight* of λ, by $m_i(\lambda)$ the *multiplicity* of i in λ, $i = 1, \dots, \infty$. Let

$$m(\lambda)! := \prod_{i=1}^{\infty} m_i(\lambda)!. \tag{1.44}$$

The partition of 0 is denoted by (0); $\ell((0)) = |(0)| = 0$. For an arbitrary sequence of indeterminates $q_1, q_2, \dots$ denote $q_\lambda := q_{\lambda_1} \cdots q_{\lambda_{\ell(\lambda)}}$; we use $\prec$ for the reverse lexicographic ordering on $\mathbb{Y}$, e.g. $\mathbb{Y}_3 = \{(3) \prec (2,1) \prec (1^4)\}$, $\mathbb{Y}_4 = \{(4) \prec (3,1) \prec (2^2) \prec (2,1^2) \prec (1^4)\}$.

There are two ways to represent the generating functions of the intersection numbers $\langle \kappa_1^{d_1} \dots \kappa_\ell^{d_\ell} \tau_{k_1} \dots \tau_{k_n} \rangle$. The first one represents the generating function of intersections of ψ- and κ-classes in terms of the generating functions of intersections of ψ-classes obtained in Theorem 1.11. The second approach is based on an explicit representation of the wave function $\psi^\kappa(z;\mathbf{t};s)$ in terms of the wave function of the Witten–Kontsevich solution. These two approaches are formulated as, respectively, the first and the second part of the following theorem.

Theorem 1.14. *Part I. For a given $\lambda \in \mathbb{Y}$ and for any $n \geq 0$*

$$\sum_{k_1,\dots,k_n \geq 0} \langle \kappa_{\lambda_1} \dots \kappa_{\lambda_{\ell(\lambda)}} \tau_{k_1} \dots \tau_{k_n} \rangle \frac{(2k_1+1)!!}{z_1^{2k_1+2}} \cdots \frac{(2k_n+1)!!}{z_n^{2k_n+2}} = (-1)^{\ell(\lambda)} \sum_{|\mu|=|\lambda|} \frac{L_{\lambda\mu}}{m(\mu)!} \operatorname*{res}_{w_1=\infty} \cdots \operatorname*{res}_{w_{\ell(\mu)}=\infty}$$
$$\frac{w_1^{2\mu_1+3}}{(2\mu_1+3)!!} \cdots \frac{w_{\ell(\mu)}^{2\mu_{\ell(\mu)}+3}}{(2\mu_{\ell(\mu)}+3)!!} F^{WK}_{\ell(\mu)+n}(w_1, \dots, w_{\ell(\mu)}, z_1, \dots, z_n)\, dw_1 \cdots dw_{\ell(\mu)}. \tag{1.45}$$

Here $L_{\lambda\mu}$ are transition matrices from the monomial basis of symmetric functions to the power sum basis [40], see Lemma 7.3 below for an explicit expression for the entries of this matrix.

Part II. The wave function corresponding to higher Weil–Petersson volume satisfies

$$\psi^{\kappa}(z;\mathbf{t};\mathbf{s}) = \exp\left(\sum_{k=1}^{\infty}\frac{h_k(-\mathbf{s})z^{2k+3}}{(2k+3)!!}\right)\sum_{\lambda\in\mathbb{Y}}\frac{(-1)^{\ell(\lambda)}s_\lambda}{m(\lambda)!}\sum_{|\mu|=|\lambda|}L_{\lambda\mu}\frac{(-1)^{\ell(\mu)}}{m(\mu)!}\partial_{t_{\mu_1+1}}\dots\partial_{t_{\mu_{\ell(\mu)}+1}}\psi^{WK}(z;\mathbf{t}). \quad (1.46)$$

Denote by $F_n^{\kappa}(z_1,\dots,z_n;\mathbf{s})$ the generating function of the higher Weil–Petersson volumes, i.e.

$$F_n^{\kappa}(z_1,\dots,z_n;\mathbf{s}) := \sum_{l\geq 0}\sum_{\substack{k_1,\dots,k_n\geq 0\\ d_1,\dots,d_l\geq 0}}\langle\kappa_1^{d_1}\dots\kappa_\ell^{d_\ell}\tau_{k_1}\dots\tau_{k_n}\rangle\frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}}\frac{s_1^{d_1}\dots s_\ell^{d_\ell}}{d_1!\cdots d_\ell!}, \quad n\geq 1. \quad (1.47)$$

Then

$$F_1^{\kappa}(z;\mathbf{s}) = \frac{1}{4\,z}\left(-A(z;\mathbf{s})\,B_z(-z;\mathbf{s}) + B_z(z;\mathbf{s})\,A(-z;\mathbf{s}) + B(z;\mathbf{s})\,A_z(-z;\mathbf{s}) - A_z(z;\mathbf{s})\,B(-z;\mathbf{s})\right), \quad (1.48)$$

$$F_n^{\kappa}(z_1,\dots,z_n;\mathbf{s}) = -\frac{1}{n}\sum_{r\in S_n}\frac{\mathrm{Tr}\left(M^{\kappa}(z_{r_1};\mathbf{s})\cdots M^{\kappa}(z_{r_n};\mathbf{s})\right)}{\prod_{j=1}^{n}(z_{r_j}^2 - z_{r_{j+1}}^2)} - \delta_{n,2}\frac{z_1^2+z_2^2}{(z_1^2-z_2^2)^2}, \quad n\geq 2. \quad (1.49)$$

Here $A(z;\mathbf{s}) := \psi^{\kappa}(z;\mathbf{0};\mathbf{s})$, $B(z;\mathbf{s}) := \psi_x^{\kappa}(z;\mathbf{0};\mathbf{s})$,

$$M^{\kappa}(z;\mathbf{s}) := \frac{1}{2}\begin{pmatrix} -\left[A(z;\mathbf{s})B(-z;\mathbf{s}) + A(-z;\mathbf{s})B(z;\mathbf{s})\right] & -2A(z;\mathbf{s})A(-z;\mathbf{s}) \\ 2B(z;\mathbf{s})B(-z;\mathbf{s}) & A(z;\mathbf{s})B(-z;\mathbf{s}) + A(-z;\mathbf{s})B(z;\mathbf{s}) \end{pmatrix} \quad (1.50)$$

(cf. definition (1.26) of the matrix Θ).

The polynomials $h_k = h_k(x_1,\dots,x_k)$, $k\geq 1$ in the formula (1.46) are defined by the well known generating function

$$1 + \sum_{k\geq 1}h_k(x)z^k = \mathrm{e}^{\sum_{i\geq 1}x_i z^i}.$$

Example 1.15. *Using Thm. 1.14 we obtain in particular*

$$\langle\kappa_{3g-3}\rangle_{g,0} = \frac{1}{24^g\cdot g!}, \qquad g\geq 1, \quad (1.51)$$

$$\langle\kappa_1\tau_{3g-3}\rangle_{g,1} = 3\frac{12g^2-12g+5}{5!!\cdot 24^g\cdot g!}, \qquad g\geq 1, \quad (1.52)$$

$$\langle\kappa_2\tau_{3g-4}\rangle_{g,1} = 3\frac{72g^3-132g^2+95g-35}{7!!\cdot 24^g\cdot g!}, \qquad g\geq 2, \quad (1.53)$$

$$\langle\kappa_3\tau_{3g-5}\rangle_{g,1} = \frac{1296g^4-3888g^3+4482g^2-2835g+945}{9!!\cdot 24^g\cdot g!}, \qquad g\geq 2. \quad (1.54)$$

Moreover, we find

$$A^{(1)}(z) = -\frac{z^5}{5!!}c + \frac{z^5}{5!!}q - \frac{z^2}{2\cdot 5!!}c, \quad (1.55)$$

$$A^{(2)}(z) = -\frac{z^7}{7!!}c + \frac{8z^7+5z}{8\cdot 7!!}q - \frac{z^4}{2\cdot 7!!}c, \tag{1.56}$$

$$A^{(1^2)}(z) = \left(\frac{z^{10}}{225} + \frac{11z^7}{1575} - \frac{z^4}{2520}\right)c + \left(-\frac{z^{10}}{225} - \frac{z^7}{210} + \frac{3z}{560}\right)q, \tag{1.57}$$

$$B^{(1)}(z) = -\frac{z^6}{5!!}q + \frac{z^3}{2\cdot 5!!}q + \frac{4z^6-6}{4\cdot 5!!}c, \tag{1.58}$$

$$B^{(2)}(z) = -\frac{z^8}{7!!}q + \frac{z^5}{2\cdot 7!!}q + \frac{8z^8-7z^2}{8\cdot 7!!}c, \tag{1.59}$$

$$B^{(1^2)}(z) = \left(-\frac{z^{11}}{225} - \frac{z^8}{210} + \frac{z^5}{150} - \frac{z^2}{240}\right)c + \left(\frac{z^{11}}{225} + \frac{4z^8}{1575} - \frac{13z^5}{2520}\right)q. \tag{1.60}$$

Here $c = c(z)$, $q = q(z)$ are Faber–Zagier series (6.10),(6.11); for a partition λ, $A^\lambda(z)$ is the coefficient of $s_\lambda := \prod_{j=1}^{\ell(\lambda)} s_{\lambda_j}$ in the **s***-expansion of $A(z;\mathbf{s})$, similarly for definition of $B^\lambda(z)$.*

Organization of the paper In Sect. 2, we briefly review the KdV hierarchy. In Sect. 3, we prove Theorem 1.2 and derive some useful lemmas. In Sect. 4, we prove Prop. 1.6 and Theorem 1.7. Application to the Witten–Kontsevich solution is presented in Sect. 6. Application to higher Weil–Petersson volumes is presented in Sect. 7. Further remarks are given in Sect. 8. We list useful formulæ and tables of intersection numbers, and apply a generalization of the Kac–Schwarz operator to higher Weil–Petersson volumes in appendices.

Acknowledgements

We would like to thank Tamara Grava, Dmitry Korotkin, Youjin Zhang, Jian Zhou for helpful discussions. We are grateful to Erik Carlsson and Fernando Rodriguez Villegas for pointing out that $L_{\lambda\mu}$ in Theorem 1.14 are transition matrices from monomial basis to homogeneous basis of symmetric functions. We thank Bertrand Eynard for bringing our attention to the papers [6], [2]. D. Y. is grateful to Youjin Zhang for his advising. Tables in Appendix C have been verified for $3g-3+n \leq 20$ with intersection numbers produced by Carel Faber's Maple program; we thank Faber for sharing it publicly. The work is partially supported by PRIN 2010-11 Grant "Geometric and analytic theory of Hamiltonian systems in finite and infinite dimensions" of Italian Ministry of Universities and Researches. M.B. is supported in part by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Fonds de recherche Nature et téchnologie du Québec (FRQNT).

2 A brief reminder of the KdV hierarchy

In this preliminary section we review some useful facts about the KdV hierarchy (1.3), and derive generalized Kac–Schwarz operators associated to string equations.

Let $\mathcal{A}_u$ be the ring of differential polynomials in u. Define a family of differential polynomials $\Omega_{p;0}(u;u_x,...,u_p)$ with zero constant term for $p \geq 0$ through the Lenard–Magri recursion

$$\partial_x \Omega_{p;0} = \frac{1}{2p+1}\left(2\,u\,\partial_x + u_x + \frac{1}{4}\partial_x^3\right)\Omega_{p-1;0}, \quad p \geq 1, \tag{2.1}$$

$$\Omega_{-1;0} = 1. \tag{2.2}$$

Here, $u_k := \partial_x^k u$, $k \geq 1$ are jet variables. Then the KdV hierarchy (1.3) can be equivalently written as

$$u_{t_p} = \partial_x \Omega_{p;0}, \quad p \geq 0. \tag{2.3}$$

The first few $\Omega_{p;0}$ are given by

$$\Omega_{0;0} = u, \tag{2.4}$$

$$\Omega_{1;0} = \frac{u^2}{2} + \frac{1}{12}u_{xx}, \tag{2.5}$$

$$\Omega_{2;0} = \frac{u^3}{6} + \left(\frac{1}{12}u\,u_{xx} + \frac{1}{24}u_x^2\right) + \frac{u_4}{240}, \tag{2.6}$$

$$\Omega_{3;0} = \frac{u^4}{24} + \left(\frac{1}{24}u^2u_{xx} + \frac{1}{24}u\,u_x^2\right) + \left(\frac{1}{240}u\,u_4 + \frac{1}{120}u_xu_3 + \frac{1}{160}u_{xx}^2\right) + \frac{u_6}{6720}. \tag{2.7}$$

In general, we have [22, 3, 18]

$$\Omega_{p;0} = \frac{1}{(2p+1)!!}Res_\partial L^{\frac{2p+1}{2}}, \quad \forall\, p \geq 0. \tag{2.8}$$

Let $\psi(z;\mathbf{t})$ and $\psi^*(z;\mathbf{t})$ be a pair of wave and dual wave functions of the KdV hierarchy. As in the Introduction, we define the following two formal series in z^{-1} by

$$\mathcal{R}(z;\mathbf{t}) := \psi(z;\mathbf{t})\,\psi^*(z;\mathbf{t}), \quad \chi(z;\mathbf{t}) := \partial_x \log\psi(z;\mathbf{t}). \tag{2.9}$$

We remind [16] that the function $\mathcal{R}$ coincides with the diagonal value of the *resolvent* of the Lax operator L. One can easily see from (1.23) and (2.8) that it gives the generating series of $\Omega_{p;0}$, i.e.

$$\mathcal{R}(z;\mathbf{t}) = 1 + \sum_{k=0}^{\infty}\frac{(2k+1)!!}{z^{2k+2}}\Omega_{k;0}(u(\mathbf{t});u_x(\mathbf{t}),\dots,u_k(\mathbf{t})). \tag{2.10}$$

It is well known that the following two lemmas hold true for $\mathcal{R}(z)$ and $\chi(z)$, for which we omit proofs.

Lemma 2.1. *The function $\mathcal{R} = \mathcal{R}(z;u;u_x,\dots)$ is the unique solution in $\mathcal{A}_u[[z^{-1}]]$ of the following ODE*

$$\mathcal{R}\,\mathcal{R}_{xx} - \frac{1}{2}\,\mathcal{R}_x^2 + (4\,u - 2\,z^2)\,\mathcal{R}^2 = -2\,z^2, \tag{2.11}$$

$$\mathcal{R} = 1 + \mathcal{O}(z^{-1}), \quad z \to \infty. \tag{2.12}$$

Furthermore, the following formula holds true for $\mathcal{R}(z)$

$$\mathcal{R}(z;u,u_x,\dots)|_{u_j=0,\,j\geq 1} = \left(1 - \frac{2\,u}{z^2}\right)^{-\frac{1}{2}}. \tag{2.13}$$

In the above formulæ it is understood that $\mathcal{R}(z;\mathbf{t}) = \mathcal{R}(z;u(\mathbf{t});u_x(\mathbf{t}),\dots)$.

One can recognize that $\mathcal{R}(z;u;u_x,\dots)$ is a full genus correction of gradient of the period of the one-dimensional Frobenius manifold [19, 22].

Lemma 2.2. *The function $\chi = \chi(z;\mathbf{t}) = \partial_x \log\psi(z;\mathbf{t})$ is the unique formal solution in $\mathcal{A}_u[[z^{-1}]]$ of the Riccati equation*

$$\chi_x + \chi^2 + 2\,u - z^2 = 0, \tag{2.14}$$

$$\chi(z;\mathbf{t}) = z + \sum_{k=1}^{\infty}\frac{\chi_k(\mathbf{t})}{z^k}, \quad z \to \infty. \tag{2.15}$$

Lemma 2.3 ([3]). *Let $\psi = \psi(z;\mathbf{t})$ and $\psi^* = \psi^*(z;\mathbf{t})$ be a pair of wave and dual wave functions of the KdV hierarchy. The following formula holds true:*

$$\psi_x\,\psi^* - \psi\,\psi^*_x = 2\,z, \tag{2.16}$$

$$\chi(z;\mathbf{t}) = \frac{1}{2}\,\partial_x \log \mathcal{R}(z;\mathbf{t}) + \frac{z}{\mathcal{R}(z;\mathbf{t})}. \tag{2.17}$$

Proof. The identity (2.16) is well-known. The proof of (2.17) is straightforward by using (2.16) and (2.9). □

3 One-point correlation functions of the KdV hierarchy

In this section we prove Theorem 1.2 by using the bilinear identities (1.6). Following [3, 17, 22], we will frequently use the following differential operator

$$\nabla(z) := \sum_{k=0}^{\infty} \frac{(2k+1)!!}{z^{2k+2}} \frac{\partial}{\partial t_k}. \tag{3.1}$$

Lemma 3.1. *For any tau-function $\tau(\mathbf{t})$ of the KdV hierarchy and any $z \in \mathbb{C}$, $z \neq 0$ we have*

$$\frac{\tau_x(\mathbf{t} - [z^{-1}])\,\tau(\mathbf{t} + [z^{-1}])}{2\,z\,\tau(\mathbf{t})^2} - \frac{\tau_x(\mathbf{t} + [z^{-1}])\,\tau(\mathbf{t} - [z^{-1}])}{2\,z\,\tau(\mathbf{t})^2} + \frac{\tau(\mathbf{t} - [z^{-1}])\,\tau(\mathbf{t} + [z^{-1}])}{\tau(\mathbf{t})^2} = 1. \tag{3.2}$$

The proof of this lemma is a straightforward application of Lemma 2.3 and hence omitted.

Proof of Theorem 1.2. Let $\tau(\mathbf{t})$ be any tau-function of the KdV hierarchy. Let $\psi(z;\mathbf{t})$ and $\psi^*(z;\mathbf{t})$ be the corresponding wave and dual wave functions (1.8). The bilinear identities (1.6), (1.14) imply that $\forall\,\mathbf{t}, \widetilde{\mathbf{t}}, w$

$$\oint_{R<|z|<|w|} \tau(\mathbf{t} - [w^{-1}] - [z^{-1}])\,\tau(\widetilde{\mathbf{t}} + [w^{-1}] + [z^{-1}]) \mathrm{e}^{\sum_{j\geq 0}\left(t_{2j+1} - \tilde{t}_{2j+1} - 2\,\frac{(2j-1)!!}{w^{2j-1}}\right)\frac{z^{2j+1}}{(2j+1)!!}}\,dz = 0. \tag{3.3}$$

Here R is a sufficiently large number. Note that

$$\mathrm{e}^{-2\sum_{j\geq 0}\frac{1}{2j+1}\left(\frac{z}{w}\right)^{2j+1}} = \frac{w-z}{w+z}. \tag{3.4}$$

Applying the operator $\nabla(w)$ to the above identity (3.3) and then setting $\widetilde{\mathbf{t}} = \mathbf{t}$, we obtain that $\forall\,\mathbf{t}, w$,

$$\sum_{k=0}^{\infty} \frac{(2k+1)!!}{w^{2k+2}} \left[\oint_{R<|z|<|w|} \left(\tau(\mathbf{t} - [w^{-1}] - [z^{-1}])\,\tau_{t_k}(\mathbf{t} + [w^{-1}] + [z^{-1}])\,\frac{w-z}{w+z}\right.\right.$$
$$\left.\left. - \frac{z^{2k+1}}{(2k+1)!!}\,\tau(\mathbf{t} - [w^{-1}] - [z^{-1}])\,\tau(\mathbf{t} + [w^{-1}] + [z^{-1}])\,\frac{w-z}{w+z}\right) dz\right] = 0. \tag{3.5}$$

Noticing now that

$$\frac{w-z}{w+z} = \frac{2\,w}{z+w} - 1, \quad z \to -w, \tag{3.6}$$

$$\frac{w-z}{w+z} = -1 + 2\,\frac{w}{z} + \mathcal{O}(z^{-2}), \quad z \to \infty \tag{3.7}$$

we have

$$\begin{aligned}
&\oint_{R<|z|<|w|} \tau(\mathbf{t}-[w^{-1}]-[z^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}]+[z^{-1}])\,\frac{w-z}{w+z}\,\frac{dz}{2i\pi}\\
&= -2\,w\,\tau(\mathbf{t})\,\tau_{t_k}(\mathbf{t}) + \oint_{R<|w|<|z|} \tau(\mathbf{t}-[w^{-1}]-[z^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}]+[z^{-1}])\,\frac{w-z}{w+z}\,\frac{dz}{2i\pi}\\
&= -2\,w\,\tau(\mathbf{t})\,\tau_{t_k}(\mathbf{t}) - \operatorname*{res}_{z=\infty}\big(\tau(\mathbf{t}-[w^{-1}]) - \tau_{t_0}(\mathbf{t}-[w^{-1}])\,z^{-1}+\mathcal{O}(z^{-2})\big)\\
&\qquad \big(\tau_{t_k}(\mathbf{t}+[w^{-1}]) + \tau_{t_0t_k}(\mathbf{t}+[w^{-1}])\,z^{-1}+\mathcal{O}(z^{-2})\big)\,\big(-1+2\,w\,z^{-1}+\mathcal{O}(z^{-2})\big)\,dz\\
&= -2\,w\,\tau(\mathbf{t})\,\tau_{t_k}(\mathbf{t}) + 2\,w\,\tau(\mathbf{t}-[w^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}])\\
&\qquad +\tau_{t_0}(\mathbf{t}-[w^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}]) - \tau(\mathbf{t}-[w^{-1}])\,\tau_{t_0t_k}(\mathbf{t}+[w^{-1}]).
\end{aligned} \tag{3.8}$$

Similarly we have

$$\begin{aligned}
&\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}}\oint_{R<|z|<|w|}\frac{z^{2k+1}}{(2k+1)!!}\,\tau(\mathbf{t}-[w^{-1}]-[z^{-1}])\,\tau(\mathbf{t}+[w^{-1}]+[z^{-1}])\,\frac{w-z}{w+z}\,\frac{dz}{2i\pi}\\
&= \oint_{R<|z|<|w|}\tau(\mathbf{t}-[w^{-1}]-[z^{-1}])\,\tau(\mathbf{t}+[w^{-1}]+[z^{-1}])\,\frac{z}{(w+z)^2}\,\frac{dz}{2i\pi}\\
&= -\tau(\mathbf{t})\,\tau(\mathbf{t}) + \tau(\mathbf{t})\,\tau(\mathbf{t}) = 0.
\end{aligned} \tag{3.9}$$

Here we have used the following expansions

$$\frac{z}{(w+z)^2} = \frac{-w}{(z+w)^2}+\frac{1}{z+w}, \quad z\to -w, \tag{3.10}$$

$$\frac{z}{(w+z)^2} = \frac{1}{z}+\mathcal{O}(z^{-2}), \quad z\to\infty. \tag{3.11}$$

Substituting (3.8) and (3.9) into (3.5) we arrive at

$$\begin{aligned}
&\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}\,\tau(\mathbf{t})^2}\bigg(\tau(\mathbf{t}-[w^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}]) + \frac{1}{2\,w}\,\tau_{t_0}(\mathbf{t}-[w^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}])\\
&\qquad -\frac{1}{2\,w}\,\tau(\mathbf{t}-[w^{-1}])\,\tau_{t_0t_k}(\mathbf{t}+[w^{-1}])\bigg) = \sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}}\frac{\tau_{t_k}(\mathbf{t})}{\tau(\mathbf{t})}.
\end{aligned} \tag{3.12}$$

Replacing $w\to -w$ in (3.12) we obtain

$$\begin{aligned}
&\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}\,\tau(\mathbf{t})^2}\bigg(\tau(\mathbf{t}+[w^{-1}])\,\tau_{t_k}(\mathbf{t}-[w^{-1}]) - \frac{1}{2\,w}\tau_{t_0}(\mathbf{t}+[w^{-1}])\,\tau_{t_k}(\mathbf{t}-[w^{-1}])\\
&\qquad +\frac{1}{2\,w}\tau(\mathbf{t}+[w^{-1}])\,\tau_{t_0t_k}(\mathbf{t}-[w^{-1}])\bigg) = \sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}}\frac{\tau_{t_k}(\mathbf{t})}{\tau(\mathbf{t})}.
\end{aligned} \tag{3.13}$$

Now let us look at the r.h.s. of (1.19). It is straightforward to compute the following products

$$\begin{aligned}
\psi_x\psi_z^* &= -\frac{\tau_{t_0}(\mathbf{t}-[z^{-1}])}{\tau(\mathbf{t})^2}\sum_{k=0}^{\infty}\tau_{t_k}(\mathbf{t}+[z^{-1}])\,\frac{(2k+1)!!}{z^{2k+2}} - \frac{\tau_{t_0}(\mathbf{t}-[z^{-1}])\tau(\mathbf{t}+[z^{-1}])}{\tau(\mathbf{t})^2}\vartheta_z\\
&\quad -z\,\frac{\psi(z;\mathbf{t})}{\tau(\mathbf{t})}\sum_{k=0}^{\infty}\tau_{t_k}(\mathbf{t}+[z^{-1}])\frac{(2k+1)!!}{z^{2k+2}}\mathrm{e}^{-\vartheta} - z\,\vartheta_z\,\psi(z;\mathbf{t})\psi^*(z;\mathbf{t})
\end{aligned}$$

$$+\frac{\psi(z;\mathbf{t})\tau_{t_0}(\mathbf{t})}{\tau(\mathbf{t})^2}\sum_{k=0}^{\infty}\tau_{t_k}(\mathbf{t}+[z^{-1}])\frac{(2k+1)!!}{z^{2k+2}}\mathrm{e}^{-\vartheta}-\frac{\tau_{t_0}(\mathbf{t})}{\tau(\mathbf{t})^2}\,\vartheta_z\,\psi(z;\mathbf{t})\psi^*(z;\mathbf{t}), \tag{3.14}$$

$$\psi_z\psi^*_x = -\psi_x\psi^*_z|_{z\to -z}, \tag{3.15}$$

$$\begin{aligned}\psi\psi^*_{xz} &= \sum_{k=0}^{\infty}\frac{(2k+1)!!}{z^{2k+2}\,\tau(\mathbf{t})^2}\,\tau(\mathbf{t}-[z^{-1}])\tau_{t_0t_k}(\mathbf{t}+[z^{-1}])-\frac{\tau(\mathbf{t}-[z^{-1}])\,\tau_{t_0}(\mathbf{t}+[z^{-1}])}{\tau(\mathbf{t})^2}\vartheta_z\\ &\quad -\psi(z;\mathbf{t})\,\psi^*(z;\mathbf{t})-\left(z\,\psi(z;\mathbf{t})+\frac{\tau_{t_0}(\mathbf{t})}{\tau(\mathbf{t})}\,\psi(z;\mathbf{t})\right)\psi^*_z(z;\mathbf{t}),\end{aligned} \tag{3.16}$$

$$\psi_{xz}\psi^* = -\psi\psi^*_{xz}|_{z\to -z}. \tag{3.17}$$

As a result we have

$$\begin{aligned}&\frac{\psi_x\psi^*_w+\psi_w\psi^*_x-\psi\psi^*_{xw}-\psi_{xw}\psi^*}{4\,w}=\\ &=\frac{1}{4\,w}\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}\tau(\mathbf{t})^2}\left[\tau(\mathbf{t}-[w^{-1}])\,\tau_{t_0t_k}(\mathbf{t}+[w^{-1}])-\tau_{t_0}(\mathbf{t}-[w^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}])\right.\\ &\qquad\left.-\tau(\mathbf{t}+[w^{-1}])\,\tau_{t_0t_k}(\mathbf{t}-[w^{-1}])+\tau_{t_0}(\mathbf{t}+[w^{-1}])\,\tau_{t_k}(\mathbf{t}-[w^{-1}])\right]\\ &\quad-\frac{1}{2}\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}\tau(\mathbf{t})^2}\left[\tau(\mathbf{t}-[w^{-1}])\,\tau_{t_k}(\mathbf{t}+[w^{-1}])+\tau(\mathbf{t}+[w^{-1}])\,\tau_{t_k}(\mathbf{t}-[w^{-1}])\right]\\ &\quad+\frac{\vartheta_w}{2\,\tau(\mathbf{t})^2}\left[\frac{1}{w}\,\tau(\mathbf{t}-[w^{-1}])\,\tau_{t_0}(\mathbf{t}+[w^{-1}])-\frac{1}{w}\,\tau(\mathbf{t}+[w^{-1}])\,\tau_{t_0}(\mathbf{t}-[w^{-1}])-2\,\tau(\mathbf{t}-[w^{-1}])\,\tau(\mathbf{t}+[w^{-1}])\right]\\ &=-\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}}\frac{\tau_{t_k}(\mathbf{t})}{\tau(\mathbf{t})}-\vartheta_w.\end{aligned} \tag{3.18}$$

The last equality uses (3.12), (3.13) and Lemma 3.1. The theorem is proved. □

Remark 3.2. *The interesting formula in Theorem 1.2 is inspired by, and can be derived alternatively from, the formulation of a Jimbo–Miwa–Ueno isomonodromic tau-function [29, 7]. We will use this formulation in a forthcoming publication for Drinfeld–Sokolov hierarchies [8].*

4 Multi-point correlation functions of the KdV hierarchy

In this section we apply Theorem 1.2 to derive generating functions of multi-point correlation functions of the KdV hierarchy. We will need the following formulae for the time derivatives of wave functions.

Lemma 4.1. *Let $\psi(z;\mathbf{t})$ and $\psi^*(z;\mathbf{t})$ be any pair of wave and dual wave functions and let $\mathcal{R}(z;\mathbf{t})=\psi(z;\mathbf{t})\,\psi^*(z;\mathbf{t})$. It follows that*

$$\nabla(z)\,\psi(w) = \frac{2\,\mathcal{R}(z)\psi_x(w)-\mathcal{R}_x(z)\psi(w)}{2\,(z^2-w^2)}, \tag{4.1}$$

$$\nabla(z)\,\psi_x(w) = \frac{\mathcal{R}_x(z)\psi_x(w)-\left[\mathcal{R}_{xx}(z)-2\,\mathcal{R}(z)(w^2-2\,u)\right]\psi(w)}{2\,(z^2-w^2)}, \tag{4.2}$$

$$\nabla(z)\,\psi_w(w) = \frac{2\,\mathcal{R}(z)\psi_{xw}(w)-\mathcal{R}_x(z)\psi_w(w)}{2\,(z^2-w^2)}-w\frac{2\,\mathcal{R}(z)\psi_x(w)-\mathcal{R}_x(z)\psi(w)}{(z^2-w^2)^2}, \tag{4.3}$$

$$\nabla(z)\,\Psi(w) \;=\; \frac{1}{2(z^2-w^2)}\begin{pmatrix} -\mathcal{R}_x(z) & -2\,\mathcal{R}(z) \\ \mathcal{R}_{xx}(z)-2\,(w^2-2\,u)\mathcal{R}(z) & \mathcal{R}_x(z)\end{pmatrix}\Psi(w), \tag{4.4}$$

$$\nabla(z)\,\Psi(w) \;=\; \frac{1}{z^2-w^2}\Theta(z)\Psi(w)+\begin{pmatrix} 0 & 0 \\ \mathcal{R}(z) & 0\end{pmatrix}\Psi(w). \tag{4.5}$$

Proof. The identity (4.1) is a standard result; e.g. see the formula (3.12) in [18] or (11.24) in [3]. The others follow from (4.1). □

Remark 4.2. *The Definition of the matrices Ψ and Θ is valid not just in a formal sense. In many cases the matrix Ψ solves a Riemann–Hilbert problem; for example if the potential $u(x)$ is in an appropriate Schwartz class or it belongs to the class of quasi-periodic functions (as in the finite gap integration problems). In these cases the identities for generating functions in the next section have also an analytic meaning and allow to tackle the problem of studying large time behaviours. This is interesting e.g. in the case of the particular solution of Witten–Kontsevich (see below), because the matrix Ψ could be written not just as a formal series but analytically in terms of Airy functions. We postpone this type of analysis to a forthcoming publication [4].*

Remark 4.3. *The matrix Θ is a resolvent of the matrix-valued Lax operator; see e.g. pages 158–159 in [16].*

4.1 Generating function of two-point correlation functions

In this subsection we prove Prop. 1.6.

Proof of Prop. 1.6. Applying the operator $\nabla(w)$ on both sides of (1.19) we obtain

$$\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}}\sum_{j=0}^{\infty}\frac{(2j+1)!!}{z^{2j+2}}\langle\langle\tau_k\tau_j\rangle\rangle \;=\frac{1}{2}\,\nabla(w)\,\mathrm{Tr}\left(\Psi^{-1}(z)\Psi_z(z)\sigma_3\right)-\nabla(w)\vartheta_z(z). \tag{4.6}$$

Clearly

$$\nabla(w)\vartheta_z(z)=\frac{z^2+w^2}{(z^2-w^2)^2}. \tag{4.7}$$

It is easy to see that $\Psi^{-1}(z)=\frac{1}{2\,z}\sigma_2\Psi(z)^T\sigma_2$. Here and below we will use Pauli matrices

$$\sigma_1=\begin{pmatrix}0&1\\1&0\end{pmatrix};\quad \sigma_2=\begin{pmatrix}0&-i\\i&0\end{pmatrix};\quad \sigma_3=\begin{pmatrix}1&0\\0&-1\end{pmatrix}. \tag{4.8}$$

We have

$$\begin{aligned}&\nabla(w)\,\mathrm{Tr}\left(\Psi^{-1}(z)\Psi_z(z)\sigma_3\right)=\\ &=\frac{1}{2\,z}\mathrm{Tr}\left[\sigma_2\nabla(w)\left(\Psi(z)^T\right)\sigma_2\Psi_z(z)\sigma_3+\sigma_2\Psi(z)^T\sigma_2\nabla(w)\left(\Psi_z(z)\right)\sigma_3\right]\\ &=\frac{2}{(w^2-z^2)^2}\mathrm{Tr}\left(\Theta(w)\Theta(z)\right)-\frac{i\,\mathrm{Tr}\left[\Psi(z)^T(\Theta(w)^T\,\sigma_2+\sigma_2\,\Theta(w))\Psi_z(z)\sigma_1\right]}{2\,z\,(w^2-z^2)}\\ &\qquad -\frac{i}{2\,z}\,\mathrm{Tr}\left[\Psi(z)^T(Q(w)^T\,\sigma_2+\sigma_2\,Q(w))\Psi_z(z)\sigma_1\right].\end{aligned} \tag{4.9}$$

Here

$$Q(w):=\begin{pmatrix}0&0\\\mathcal{R}(w)&0\end{pmatrix}. \tag{4.10}$$

Noticing that $Q(w)^T\,\sigma_2+\sigma_2\,Q(w)=0,\ \Theta(w)^T\,\sigma_2+\sigma_2\,\Theta(w)=0$, we have

$$\nabla(w)\left[\mathrm{Tr}\left(\Psi^{-1}(z)\Psi_z(z)\sigma_3\right)\right]=\frac{2}{(w^2-z^2)^2}\mathrm{Tr}\left(\Theta(w)\Theta(z)\right). \tag{4.11}$$

Hence

$$\sum_{k=0}^{\infty}\frac{(2k+1)!!}{w^{2k+2}}\sum_{j=0}^{\infty}\frac{(2j+1)!!}{z^{2j+2}}\langle\langle\tau_k\tau_j\rangle\rangle=\frac{\mathrm{Tr}\left(\Theta(w)\Theta(z)\right)-z^2-w^2}{(w^2-z^2)^2} \tag{4.12}$$

$$=\frac{\frac{1}{2}\mathcal{R}_x(w)\mathcal{R}_x(z)-\frac{1}{2}(\mathcal{R}_{xx}(w)\mathcal{R}(z)+\mathcal{R}_{xx}(z)\mathcal{R}(w))+(z^2+w^2-4\,u)\mathcal{R}(w)\mathcal{R}(z)-z^2-w^2}{(w^2-z^2)^2}. \tag{4.13}$$

Substituting (2.11) into the formula we obtain (1.24). The theorem is proved. □

Example 4.4. *The first few two-point correlation functions are given by*

$$\langle\langle\tau_0^2\rangle\rangle=u,\quad \langle\langle\tau_0\tau_1\rangle\rangle=\frac{u^2}{2}+\frac{1}{12}u_{xx}, \tag{4.14}$$

$$\langle\langle\tau_1^2\rangle\rangle=\frac{u^3}{3}+\left(\frac{u_x^2}{24}+\frac{u\,u_{xx}}{6}\right)+\frac{1}{144}u_{xxxx}, \tag{4.15}$$

$$\langle\langle\tau_1\tau_2\rangle\rangle=\frac{u^4}{8}+\left(\frac{u\,u_x^2}{12}+\frac{1}{8}u^2u_{xx}\right)+\left(\frac{1}{90}u\,u_4+\frac{23\,u_{xx}^2}{1440}+\frac{1}{60}\,u_xu_3\right)+\frac{1}{2880}u_6. \tag{4.16}$$

To end this subsection we point out that, for an arbitrary solution $u(\mathbf{t})$ of the KdV hierarchy, the Sato tau-function τ and the axiomatic tau-function τ^{KdV} defined in [22] coincide up to a gauge factor of the form $\mathrm{e}^{\alpha_{-1}+\sum_{j\geq 0}t_j\alpha_j}$. Indeed, by definition we know that $\forall\, p,q,r$,

$$\langle\langle\tau_p\tau_q\rangle\rangle=\langle\langle\tau_q\tau_p\rangle\rangle,\quad \langle\langle\tau_p\tau_q\tau_r\rangle\rangle=\langle\langle\tau_q\tau_r\tau_p\rangle\rangle=\langle\langle\tau_r\tau_p\tau_q\rangle\rangle. \tag{4.17}$$

From Theorem 1.6 and Lemma 2.1 we know that $\langle\langle\tau_p\tau_q\rangle\rangle\in\mathcal{A}_u$ and that

$$\langle\langle\tau_p\tau_q\rangle\rangle|_{u_j=0,\,j\geq 1}=\frac{u^{p+q+1}}{p!q!(p+q+1)}. \tag{4.18}$$

So τ satisfies the defining properties of τ^{KdV}.

4.2 Generating functions of multi-point correlation functions

Notation. For a permutation $r\in S_n$, $n\geq 2$ denote

$$P(r)=-\prod_{j=1}^{n}\frac{1}{z_{r_j}^2-z_{r_{j+1}}^2} \tag{4.19}$$

where r_{n+1} is understood as r_1.

Lemma 4.5. *The following formula holds true:*

$$\nabla(z)\Theta(w)=\frac{1}{z^2-w^2}[\Theta(z),\Theta(w)]+[Q(z),\Theta(w)]. \tag{4.20}$$

with Q defined in (4.10).

Proof. We have, using the Leibniz rule,

$$\begin{aligned}
\nabla(z)\,\Theta(w) &= \nabla(z)\left[\Psi(w)\sigma_3\Psi(w)^{-1}\right] = \nabla(z)\left(\Psi(w)\right)\sigma_3\Psi(w)^{-1} + \Psi(w)\sigma_3\nabla(z)\left(\Psi(w)^{-1}\right) \\
&= \left[\frac{1}{z^2-w^2}\Theta(z)\Psi(w) + Q(z)\Psi(w)\right]\sigma_3\Psi(w)^{-1} \\
&\qquad + \Psi(w)\sigma_3\left[\frac{1}{z^2-w^2}\Psi(w)^{-1}\sigma_2\Theta(z)^T\sigma_2 + \Psi(w)^{-1}\sigma_2 Q(z)^T\sigma_2\right] \\
&= \frac{1}{z^2-w^2}[\Theta(z),\Theta(w)] + [Q(z),\Theta(w)]. \qquad (4.21)
\end{aligned}$$

□

Proof of Theorem 1.7. We use mathematical induction on n. For $n=2$, the theorem has been already verified in Prop. 1.6. Suppose it is true for $n=p\,(p\geq 2)$. For $n=p+1$,

$$\begin{aligned}
&\sum_{k_1,\ldots,k_{p+1}=0}^{\infty}\frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_{p+1}+1)!!}{z_{p+1}^{2k_{p+1}+2}}\langle\langle\tau_{k_1}\ldots\tau_{k_{p+1}}\rangle\rangle \\
&= \nabla(z_{p+1})\sum_{k_1,\ldots,k_p=0}^{\infty}\frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_p+1)!!}{z_{p+1}^{2k_p+2}}\langle\langle\tau_{k_1}\ldots\tau_{k_p}\rangle\rangle \\
&= \nabla(z_{p+1})\left(\frac{1}{p}\sum_{r\in S_p}P(r)\,\mathrm{Tr}\left(\Theta(z_{r_1})\cdots\Theta(z_{r_p})\right) - \delta_{p,2}\frac{z_1^2+z_2^2}{(z_1^2-z_2^2)^2}\right) \\
&= \frac{1}{p}\sum_{r\in S_p}P(r)\sum_{j=1}^{p}\mathrm{Tr}\left[\Theta(z_{r_1})\cdots\nabla(z_{p+1})\big(\Theta(z_{r_j})\big)\cdots\Theta(z_{r_p})\right] \\
&= \frac{1}{p}\sum_{r\in S_p}P(r)\sum_{j=1}^{p}\mathrm{Tr}\left[\Theta(z_{r_1})\cdots\left(\frac{1}{z_{p+1}^2-z_{r_j}^2}[\Theta(z_{p+1}),\Theta(z_{r_j})] + [Q(z_{p+1}),\Theta(z_{r_j})]\right)\cdots\Theta(z_{r_p})\right] \\
&\overset{\star}{=} \frac{1}{p}\sum_{r\in S_p}P(r)\sum_{j=1}^{p}\left(\frac{1}{z_{p+1}^2-z_{r_j}^2} - \frac{1}{z_{p+1}^2-z_{r_{j-1}}^2}\right)\mathrm{Tr}\left(\Theta(z_{r_1})\cdots\Theta(z_{r_{j-1}})\Theta(z_{p+1})\Theta(z_{r_j})\cdots\Theta(z_{r_p})\right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{p}\sum_{j=1}^{p}\sum_{r\in S_p}P(r)\frac{z_{r_j}^2-z_{r_{j-1}}^2}{(z_{p+1}^2-z_{r_j}^2)(z_{p+1}^2-z_{r_{j-1}}^2)}\mathrm{Tr}\left(\Theta(z_{p+1})\Theta(z_{r_j})\cdots\Theta(z_{r_p})\Theta(z_{r_1})\cdots\Theta(z_{r_{j-1}})\right) \\
&= \frac{1}{p}\sum_{j=1}^{p}\sum_{r\in S_p}P\left([p+1,r_j,\ldots,r_p,r_1,\ldots,r_{j-1}]\right)\mathrm{Tr}\left(\Theta(z_{p+1})\Theta(z_{r_j})\cdots\Theta(z_{r_p})\Theta(z_{r_1})\cdots\Theta(z_{r_{j-1}})\right) \\
&\overset{\dagger}{=} \sum_{r\in S_p}P\left([p+1,r]\right)\,\mathrm{Tr}\left(\Theta(z_{p+1})\Theta(z_{r_1})\cdots\Theta(z_{r_p})\right) \\
&= \frac{1}{p+1}\sum_{r\in S_{p+1}}P(r)\,\mathrm{Tr}\left(\Theta(z_{r_1})\cdots\Theta(z_{r_{p+1}})\right). \qquad (4.22)
\end{aligned}$$

Here, for a permutation $r=[r_1,\ldots,r_\ell]\in S_\ell$, $r_0:=r_\ell$, and we have used the facts that both $P(r)$ and matrix trace of products are invariant under the cyclic permutation. In the step marked with an asterisk, we have used

that the terms of the form

$$\sum_{j=1}^{p} \mathrm{Tr}\Big[\Theta(z_{r_1})\cdots\overbrace{[Q(z_{p+1}),\Theta(z_{r_j})]}^{j\text{–th place}}\cdots\Theta(z_{r_p})\Big] \tag{4.23}$$

are zero; to see this the reader can notice that the above is the infinitesimal action of the conjugation action by $\mathrm{e}^{sQ(z_{p+1})}$ and the trace $\mathrm{Tr}(\Theta(z_{r_1})\cdots\Theta(z_{r_p}))$ is invariant under simultaneous conjugation of the matrices. In the step marked with † we have relabeled the summation over permutations. In the last step we have used that the summand is invariant under cyclic permutations of the indices. The theorem is proved. □

5 On string equation and Kac–Schwarz operator

In the remaining part of this paper we will consider a subset of solutions of the KdV hierarchy [22] whose tau-functions are specified by string equations:

$$L_{-1}\tau = 0, \qquad L_{-1} := \sum_{k\geq 0} \tilde{t}_{k+1}\frac{\partial}{\partial t_k} + \frac{1}{2}\tilde{t}_0^2 \tag{5.1}$$

where $\tilde{t}_k = t_k - c_k,\ k \geq 0$, with arbitrary constants c_k. Note that the string equation (5.1) uniquely determines a tau-function up to a multiplicative constant independent of $\mathbf{t}$. The Witten–Kontsevich tau-function as well as the generating function of higher Weil–Petersson volumes are particular examples in this class of solutions. E.g., for the Witten–Kontsevich solution $c_1 = 1$, $c_k = 0$ for $k \neq 1$.

It was observed by V. Kac and A.S. Schwarz [31] that the string equation for the Witten–Kontsevich tau-function implies a differential equation in the spectral parameter z for the associated wave function. The equation can be written in terms of the linear differential operator $S_z = \frac{1}{z}\partial_z - \frac{1}{2z^2} - z$ that will be called *Kac–Schwarz operator*. In this section, following A. Buryak [12], we give a generalization of the Kac–Schwarz operator for the general class of solutions satisfying (5.1).

Definition 5.1. *The generalized Kac–Schwarz operator S_z associated to the string equation (5.1) is defined as the following differential operator of z :*

$$S_z = \frac{1}{z}\partial_z - \frac{1}{2z^2} - \sum_{k=0}^{\infty}\frac{c_k}{(2k-1)!!}z^{2k-1}. \tag{5.2}$$

where the constants c_k are the shifts in the times t_k in (5.1).

The above definition also can been generalized to the Gelfand–Dickey hierarchy in [31, 9].

Proposition 5.2. *Let $K(z;\mathbf{t}) := \psi(z;\mathbf{t})\cdot\tau(\mathbf{t}) = \tau(\mathbf{t} - [z^{-1}])\,\mathrm{e}^{\vartheta(z)}$. Then the following formula holds true:*

$$L_{-1}K = S_z K. \tag{5.3}$$

Proof. On one hand we have

$$\begin{aligned} L_{-1}K &= L_{-1}\left(\tau(\mathbf{t}-[z^{-1}])\cdot\mathrm{e}^{\vartheta}\right) \\ &= \mathrm{e}^{\vartheta}\cdot\sum_{k=0}^{\infty}\tilde{t}_{k+1}\frac{\partial\tau(\mathbf{t}-[z^{-1}])}{\partial t_k} + \sum_{k\geq 0}\frac{\tilde{t}_{k+1}\,z^{2k+1}}{(2k+1)!!}K + \frac{(\tilde{t}_0)^2}{2}K \end{aligned}$$

$$
\begin{aligned}
&= \mathrm{e}^{\vartheta}\cdot\sum_{k=0}^{\infty}\left(\tilde{t}_{k+1}-\frac{(2k+1)!!}{z^{2k+3}}\right)\frac{\partial\tau(\mathbf{t}-[z^{-1}])}{\partial t_k}+\sum_{k\geq 0}\frac{\tilde{t}_{k+1}\,z^{2k+1}}{(2k+1)!!}K+\frac{(\tilde{t}_0-z^{-1})^2}{2}K\\
&\quad+\mathrm{e}^{\vartheta}\cdot\sum_{k=0}^{\infty}\frac{(2k+1)!!}{z^{2k+3}}\frac{\partial\tau(\mathbf{t}-[z^{-1}])}{\partial t_k}-\frac{1}{2z^2}K+\frac{1}{z}\tilde{t}_0K\\
&= \mathrm{e}^{\vartheta}\cdot\sum_{k=0}^{\infty}\frac{(2k+1)!!}{z^{2k+3}}\frac{\partial\tau(\mathbf{t}-[z^{-1}])}{\partial t_k}-\frac{1}{2z^2}K+\sum_{k\geq -1}\frac{\tilde{t}_{k+1}\,z^{2k+1}}{(2k+1)!!}K. \qquad (5.4)
\end{aligned}
$$

On another hand we have

$$
\frac{1}{z}\partial_z K=\mathrm{e}^{\vartheta}\sum_{k=0}^{\infty}\frac{(2k+1)!!}{z^{2k+3}}\frac{\partial\tau(\mathbf{t}-[z^{-1}])}{\partial t_k}+\sum_{k\geq 0}\frac{t_k z^{2k-1}}{(2k-1)!!}K \qquad (5.5)
$$

Since $\widetilde{t}_k=t_k-c_k$, we obtain $L_{-1}K=\frac{1}{z}\partial_z K-\frac{1}{2z^2}K-\sum_{k=0}^{\infty}\frac{c_k z^{2k-1}}{(2k-1)!!}K$, which was our statement. □

Proposition 5.3. *Let $u_0(x):=\partial_x^2\log\tau(x,\mathbf{0})$ and let $f(z;x):=\psi(z;x,\mathbf{0})$; then*

$$
S_z f(z;x)=-c_1 f_x(z;x)-\sum_{k\geq 1}c_{k+1}\psi_{t_k}(z;x,\mathbf{0}), \qquad (5.6)
$$

$$
S_z f_x(z;x)=-c_1\left(z^2-2u_0(x)\right)f(z;x)-\sum_{k\geq 1}c_{k+1}\psi_{t_0t_k}(z;x,\mathbf{0}), \qquad (5.7)
$$

$$
\psi_{t_k}(z;x,\mathbf{0})=\frac{1}{(2k+1)!!}\sum_{i=0}^{k}(2i-1)!!\,z^{2k-2i}\left(\Omega_{i-1;0}|_{u\to u_0}f_x(z;x)-\frac{1}{2}f(z;x)\partial_x\Omega_{i-1;0}|_{u\to u_0}\right), \qquad (5.8)
$$

$$
\begin{aligned}
\psi_{t_0t_k}(z;x,\mathbf{0})=\frac{1}{(2k+1)!!}\sum_{i=0}^{k}(2i-1)!!\,z^{2k-2i}&\left(\partial_x\Omega_{i-1;0}|_{u\to u_0}f_x(z;x)+\Omega_{i-1;0}|_{u\to u_0}\left(z^2-2u_0(x)\right)f(z;x)\right.\\
&\left.-\frac{1}{2}f_x(z;x)\partial_x\Omega_{i-1;0}|_{u\to u_0}-\frac{1}{2}f(z;x)\partial_x^2\Omega_{i-1;0}|_{u\to u_0}\right). \qquad (5.9)
\end{aligned}
$$

Proof. Taking $t_k=0$, $k\geq 1$ in (5.3) gives

$$
S_z f(z;x)=\frac{c_0^2}{2}f(z;x)-\sum_{k\geq 0}c_{k+1}\frac{\partial\psi}{\partial t_k}(z;x,\mathbf{0})-\sum_{k\geq 0}c_{k+1}\frac{\partial\log\tau}{\partial t_k}(x,\mathbf{0})f(z;x). \qquad (5.10)
$$

Taking $t_k=0$, $k\geq 1$ in (5.1) and substituting it into the above equality one obtains (5.6). Equation (5.7) is obtained by taking x-derivative of (5.6). Equation (5.8) can be derived from the defining equations (1.11) and (1.12), which is a standard expression hence omitted. Taking the x-derivative in (5.8) yields (5.9). □

We will apply in Appendix B results of Propositions 5.2, 5.3 to computation of higher Weil–Petersson volumes.

6 Application to the Witten–Kontsevich tau-function

In this section we consider a particular solution of the KdV hierarchy (1.3). Namely, we are going to apply Theorem 1.2, Theorem 1.7 and the Witten-Kontsevich theorem to derive n-point functions of the intersection numbers (1.31).

To do so we first derive the *initial datum* of the wave function $\psi^{WK}(z;\mathbf{t})$. Denote $f^{WK}(z;x)=\psi^{WK}(z;\mathbf{t})|_{\mathbf{t}_{\geq 1}=\mathbf{0}}$. Noting that $u_0^{WK}(x)=x$, we have

$$f^{WK}_{xx}+2\,xf^{WK}=z^2f^{WK}. \tag{6.1}$$

Solutions of this ODE can be expressed as linear combinations of Airy functions. Imposing the asymptotic condition $f^{WK}(z;x)=\big(1+\mathcal{O}(z^{-1})\big)\exp(xz)$, it is easy to obtain

$$f^{WK}(z;x)=G(z)\sqrt{2\pi z}\,\mathrm{e}^{\frac{z^3}{3}}2^{\frac{1}{3}}\mathrm{Ai}\left(2^{-\frac{2}{3}}(z^2-2\,x)\right) \tag{6.2}$$

where $G(z)=1+\mathcal{O}(z^{-1})$ is yet to be determined and is a function independent of x. The asymptotic expansion of (6.2) is of the indicated form only in a suitable sector; the dual wave function admitting the required asymptotic in the same sector is found in Appendix A. To fix the gauge freedom we need to enforce the *string equation*

$$\sum_{k=0}^{\infty}t_{k+1}\frac{\partial Z}{\partial t_k}+\frac{t_1^2}{2}Z=\frac{\partial Z}{\partial t_1}. \tag{6.3}$$

which is derived by Witten [52]. One immediately reads off from (6.3) that

$$c_0=0,\ c_1=1,\ c_k=0\,(k\geq 2). \tag{6.4}$$

So, due to Definition 5.1, the Kac–Schwarz operator associated to $Z(\mathbf{t})$ reads

$$S_z^{WK}=\frac{1}{z}\partial_z-\frac{1}{2z^2}-z. \tag{6.5}$$

Prop. 5.3 with the values (6.4) yields immediately that

$$S_z^{WK}f^{WK}(z;x)=-\partial_x f^{WK}(z;x)=\sqrt{2\pi z}\,\mathrm{e}^{\frac{z^3}{3}}2^{\frac{2}{3}}G(z)\mathrm{Ai}'\left(2^{-\frac{2}{3}}(z^2-2\,x)\right). \tag{6.6}$$

On the other hand from (6.2) we have

$$S_z^{WK}f^{WK}(z;x)=\sqrt{2\pi z}\,\mathrm{e}^{\frac{z^3}{3}}2^{\frac{1}{3}}\left(2^{\frac{1}{3}}G(z)\mathrm{Ai}'\left(2^{-\frac{2}{3}}(z^2-2\,x)\right)+\frac{G'(z)}{z}\mathrm{Ai}\left(2^{-\frac{2}{3}}(z^2-2\,x)\right)\right). \tag{6.7}$$

Comparing (6.6), (6.7) we conclude that $G(z)$ is a constant and thus $G(z)\equiv 1$.

Definition 6.1. *We define* $A^{WK}(z):=\psi^{WK}(z;\mathbf{0})=f^{WK}(z;0),\quad B^{WK}(z):=\psi^{WK}_x(z;\mathbf{0})=f^{WK}_x(z;0)$.

Setting $x=0$ in (6.7) yields

$$A^{WK}(z)=\sqrt{2\pi z}\,\mathrm{e}^{\frac{z^3}{3}}2^{\frac{1}{3}}\mathrm{Ai}\left(2^{-\frac{2}{3}}z^2\right)\simeq c(z),\quad B^{WK}(z)=-\sqrt{2\pi z}\,\mathrm{e}^{\frac{z^3}{3}}2^{\frac{2}{3}}\,\mathrm{Ai}'\left(2^{-\frac{2}{3}}z^2\right)\simeq z\,q(z) \tag{6.8}$$

where the symbol $\simeq$ stands for asymptotic equivalence, and $c(z)$ and $q(z)$ are known as the Faber–Zagier series [28, 12] which can be computed from the known expansions of Airy functions

$$\mathrm{Ai}(z)\sim\frac{\mathrm{e}^{-\zeta}}{2\sqrt{\pi}z^{1/4}}\sum_{k=0}^{\infty}\frac{(6k-1)!!}{(2k-1)!!}\frac{(-216\,\zeta)^{-k}}{k!},\quad \zeta=\frac{2}{3}z^{3/2},\quad z\to\infty,\quad |\arg z|<\pi. \tag{6.9}$$

So

$$c(z)=\sum_{k=0}^{\infty}C_k\,z^{-3k},\quad C_k=\frac{(-1)^k}{288^k}\frac{(6k)!}{(3k)!(2k)!}, \tag{6.10}$$

$$q(z)=\sum_{k=0}^{\infty}q_k\,z^{-3k},\quad q_k=\frac{1+6k}{1-6k}C_k. \tag{6.11}$$

For future reference, we also have, from Prop. 5.3

Lemma 6.2. *$A^{WK}(z), B^{WK}(z)$ satisfy the following system of ODEs*

$$\begin{cases} S_z^{WK} A^{WK}(z) = -B^{WK}(z), \\ S_z^{WK} B^{WK}(z) = -z^2 A^{WK}(z). \end{cases} \tag{6.12}$$

The Faber–Zagier series satisfy the well-known relation

$$c(z)\, q(-z) + c(-z)\, q(z) = 2, \tag{6.13}$$

which expresses (2.16) of Lemma 2.3. Furthermore they possess the following properties.

Lemma 6.3. *The following equalities hold true:*

$$c(z) \cdot c(-z) = \sum_{g=0}^{\infty} \frac{(6g-1)!!}{24^g \cdot g!} z^{-6g}, \tag{6.14}$$

$$q(z) \cdot q(-z) = -\sum_{g=0}^{\infty} \frac{6g+1}{6g-1} \frac{(6g-1)!!}{24^g \cdot g!} z^{-6g}, \tag{6.15}$$

$$c(z) \cdot q(-z) = 1 - \frac{1}{2} \sum_{g=1}^{\infty} \frac{(6g-5)!!}{24^{g-1} \cdot (g-1)!} z^{-6g+3}, \tag{6.16}$$

$$q(z) \cdot c(-z) = 1 + \frac{1}{2} \sum_{g=1}^{\infty} \frac{(6g-5)!!}{24^{g-1} \cdot (g-1)!} z^{-6g+3}. \tag{6.17}$$

Proof. By straightforward computations and by using summation formulæ for hypergeometric series (i.e. Dixon's identities and contiguous identities [35]). □

Due to the definition of $Z(\mathbf{t})$, the relation between the intersection numbers and the n-point correlation functions of $Z(\mathbf{t})$ consists simply in an evaluation at $\mathbf{t} = 0$:

$$\langle \tau_{k_1} \dots \tau_{k_n} \rangle = \langle\langle \tau_{k_1} \dots \tau_{k_n} \rangle\rangle|_{\mathbf{t}=0}. \tag{6.18}$$

Proof of Theorem 1.11. Let us first consider one-point intersection numbers. According to Theorem 1.2 and the Witten-Kontsevich theorem, we have

$$\begin{aligned} F_1^{WK}(z) &= \frac{1}{4}\left[q(z)\, c'(-z) + c'(z)\, q(-z) + \frac{c(-z)}{z}\left(q(z) + z\, q'(z)\right) + \frac{c(z)}{z}\left(-q(-z) + z\, q'(-z)\right)\right] \\ &= z^2 \left(1 - \frac{c(z)c(-z) + q(z)q(-z)}{2}\right) = \sum_{g=1}^{\infty} \frac{(6g-3)!!}{24^g \cdot g!} z^{-(6g-2)}. \end{aligned} \tag{6.19}$$

Thus we obtain

$$\langle \tau_j \rangle = \begin{cases} \frac{1}{24^g \cdot g!} & j = 3g-2, \\ 0 & \text{otherwise}, \end{cases} \tag{6.20}$$

i.e. $\langle \tau_{3g-2} \rangle = \langle \tau_{3g-2} \rangle_g = \frac{1}{24^g \cdot g!}$.

Now we consider the n-point function for intersection numbers of ψ-classes. Substituting (6.8) into (1.27) and using Lemma 6.3 we obtain

$$\begin{aligned}\Theta(z;0) &= \frac{1}{2}\begin{pmatrix} z\,(c(z)q(-z)-c(-z)q(z)) & -2\,c(z)c(-z) \\ -2z^2q(z)q(-z) & -z\,(c(z)q(-z)-c(-z)q(z)) \end{pmatrix} \\ &= \frac{1}{2}\begin{pmatrix} -\sum_{g=1}^{\infty}\frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!}z^{-6g+4} & -2\sum_{g=0}^{\infty}\frac{(6g-1)!!}{24^g\cdot g!}z^{-6g} \\ 2\sum_{g=0}^{\infty}\frac{6g+1}{6g-1}\frac{(6g-1)!!}{24^g\cdot g!}z^{-6g+2} & \sum_{g=1}^{\infty}\frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!}z^{-6g+4} \end{pmatrix} =: M(z).\end{aligned} \tag{6.21}$$

So we have

$$F_n^{WK}(z_1,\dots,z_n) = \frac{1}{n}\sum_{r\in S_n} P(r)\,\mathrm{Tr}\,(M(z_{r_1})\cdots M(z_{r_n})) - \delta_{n,2}\frac{z_1^2+z_2^2}{(z_1^2-z_2^2)^2}. \tag{6.22}$$

The theorem is proved. □

Theorem 1.11 allows a very efficient way of computing the intersection numbers of ψ-classes. We list in Appendix C several of them. The initial values of the functions $\mathcal{R}(z;\mathbf{t})$, $\mathcal{R}_x(z;\mathbf{t})$, $\mathcal{R}_{xx}(z;\mathbf{t})$ are immediately read off from Thm 1.11.

Corollary 6.4. *For the Witten–Kontsevich solution of the KdV hierarchy we have*

$$\mathcal{R}(z;\mathbf{0}) = \sum_{g=0}^{\infty}\frac{(6g-1)!!}{24^g\cdot g!}z^{-6g}, \quad \mathcal{R}_x(z;\mathbf{0}) = \sum_{g=1}^{\infty}\frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!}z^{-6g+4}, \tag{6.23}$$

$$\mathcal{R}_{xx}(z;\mathbf{0}) = \sum_{g=1}^{\infty}\frac{(6g-3)!!}{24^{g-1}\cdot(g-1)!}z^{-6g+2}. \tag{6.24}$$

Relationship with topological recursion. Let us briefly comment on the relationship between the formulæ (1.36) and functions of Eynard–Orantin type, sometimes referred to as solutions of the *topological recursions*. Recall that $Z(\mathbf{t})$ admits the genus expansion

$$\log Z(\mathbf{t}) = \sum_{g=0}^{\infty}\mathcal{F}_g(\mathbf{t}), \tag{6.25}$$

where $\mathcal{F}_g$ are genus g free energies corresponding to the Witten–Kontsevich solution

$$\mathcal{F}_g(\mathbf{t}) = \sum_{n=0}^{\infty}\frac{1}{n!}\sum_{k_1+\ldots+k_n=3g-3+n}\langle\tau_{k_1}\cdots\tau_{k_n}\rangle_{g,n}\,t_{k_1}\cdots t_{k_n}. \tag{6.26}$$

Define

$$W_{g,n}^{WK}(z_1,\dots,z_n) = \nabla(z_1)\cdots\nabla(z_n)\mathcal{F}_g(\mathbf{t})\,|_{\mathbf{t}=\mathbf{0}} \tag{6.27}$$

where we remind the reader that

$$\nabla(z) = \sum_{k=0}^{\infty}\frac{(2k+1)!!}{z^{2k+2}}\frac{\partial}{\partial t_k}. \tag{6.28}$$

Hence we have

$$F_n^{WK}(z_1,\dots,z_n)=\sum_{g=0}^{\infty}W_{g,n}^{WK}(z_1,\dots,z_n). \tag{6.29}$$

The functions $W_{g,n}^{WK}(z_1,\dots,z_n)$ are introduced by Eynard and Orantin [23]; see also [53, 54]. It is proved for example in [54] that Eynard–Orantin's topological recursions for $W_{g,n}^{WK}$ are equivalent to the Virasoro constraints [17, 22].

7 Application to higher Weil–Petersson volumes

In this section we apply Theorem 1.7 to the solution of higher Weil–Petersson volumes, for which we mean intersection numbers of mixed κ- and ψ-classes:

$$\langle\kappa_1^{d_1}\dots\kappa_\ell^{d_\ell}\tau_{k_1}\dots\tau_{k_n}\rangle_{g,n}:=\int_{\overline{\mathcal{M}}_{g,n}}\psi_1^{k_1}\dots\psi_n^{k_n}\kappa_1^{d_1}\dots\kappa_\ell^{d_\ell}. \tag{7.1}$$

These numbers are zero unless

$$\sum_{j=1}^{\ell}j\,d_j+\sum_{j=1}^{n}k_j=3g-3+n. \tag{7.2}$$

Theorem 7.1 ([32, 22, 46, 38])**.** *The partition function $Z^\kappa(\mathbf{t};\mathbf{s})$ defined by*

$$Z^\kappa(\mathbf{t};\mathbf{s})=\exp\left(\sum_{g=0}^{\infty}\sum_{n,\ell\geq 0}\frac{1}{n!}\sum_{\sum jd_j+\sum k_j=3g-3+n}\langle\kappa_1^{d_1}\dots\kappa_\ell^{d_\ell}\tau_{k_1}\dots\tau_{k_n}\rangle_{g,n}\,t_{k_1}\cdots t_{k_n}\frac{s_1^{d_1}\dots s_\ell^{d_\ell}}{d_1!\cdots d_\ell!}\right) \tag{7.3}$$

is a particular tau-function of the KdV hierarchy (1.3). Moreover,

$$Z^\kappa(\mathbf{t};\mathbf{s})=Z(t_0,t_1,t_2-h_1(-\mathbf{s}),t_3-h_2(-\mathbf{s}),\dots,t_{k+1}-h_k(-\mathbf{s}),\dots) \tag{7.4}$$

where $h_k(\mathbf{s})$ are polynomials in $s_1,s_2,\dots$ defined through

$$\sum_{k=0}^{\infty}h_k(\mathbf{s})\,x^k=\exp\left(\sum_{j=1}^{\infty}s_jx^j\right). \tag{7.5}$$

Observe that, equivalently $Z^\kappa(\mathbf{t};\mathbf{s})$ has the form

$$Z^\kappa(\mathbf{t};\mathbf{s})=\exp\left(\sum_{g=0}^{\infty}\sum_{n,q,\ell\geq 0}\sum_{\substack{\sum m_j=n\\ \sum jd_j+\sum jm_j=3g-3+n}}\langle\kappa_1^{d_1}\dots\kappa_\ell^{d_\ell}\tau_0^{m_0}\dots\tau_q^{m_q}\rangle_{g,n}\,\frac{t_0^{m_0}\cdots t_q^{m_q}}{m_0!\cdots m_q!}\frac{s_1^{d_1}\dots s_\ell^{d_\ell}}{d_1!\cdots d_\ell!}\right). \tag{7.6}$$

It is easy to see that $Z(\mathbf{t})=Z^\kappa(\mathbf{t};\mathbf{0})$.

There exist several methods for computing the integrals (7.1), including application of the Virasoro constraints [43, 44, 24, 46, 54, 38], the *quasi-triviality* approach [22, 57, 28], as well as an interesting method in the original paper of [1]. We propose a yet different approach, based on Thm. 1.7 and Thm. 7.1.

7.1 Proof of Theorem 1.14

Before entering into the proof proper, we need a few preparations.

Lemma 7.2. *The following formula holds true*

$$\exp\left(\sum_{m\geq 1} h_m(\mathbf{s})q_m\right) = \sum_{\lambda\in\mathbb{Y}} \frac{s_\lambda}{m(\lambda)!} \sum_{|\mu|=|\lambda|} L_{\lambda\mu}\frac{q_\mu}{m(\mu)!}, \tag{7.7}$$

where $L_{\lambda\mu}$ is the transition matrix from the monomial basis to the power sum basis. We have used the shorthands

$$s_\lambda := \prod_{j=1}^{\ell(\lambda)} s_{\lambda_j}\ , \qquad q_\mu := \prod_{j=1}^{\ell(\mu)} q_{\mu_j}\ . \tag{7.8}$$

Proof. By Taylor expansion of the exponential we have

$$\begin{aligned}\exp\left(\sum_{m\geq 1} h_m(\mathbf{s})q_m\right) &= \sum_{n=0}^{\infty}\sum_{\sum k_j=n}\prod_{j=1}^{\infty}\frac{q_j^{k_j}}{k_j!}\prod_{j=1}^{\infty}h_j(\mathbf{s})^{k_j}\\ &= \sum_{\mu\in\mathbb{Y}}\frac{q_\mu}{m(\mu)!}h_\mu(\mathbf{s}) = \sum_{\mu\in\mathbb{Y}}\frac{q_\mu}{m(\mu)!}\sum_{|\lambda|=|\mu|}L_{\lambda\mu}s_\lambda,\end{aligned} \tag{7.9}$$

where the last equality uses the definition of the transition matrix between homogeneous basis and power sum basis. □

It is known that the matrix $(L_{\lambda\mu})$ equals to the product of the character table and the Kostka matrix. Now we are going to give two explicit formulæ for $L_{\lambda\mu}$ for given λ,μ. Write $\mathbb{Y} = \{\sigma_0 \prec \sigma_1 \prec \sigma_2 \prec \ldots\}$, where $\sigma_0 = (0)$, $\sigma_1 = (1)$, $\sigma_2 = (2)$, $\sigma_3 = (1^2)$, etc. Let $p(n)$ be the number of partitions of weight n, i.e. $p(0) = p(1) = 1$, $p(2) = 2$, $p(3) = 3, \ldots$; and let $r_n := \sum_{k=1}^{n} p(k)$, $n \geq 1$, $r_0 := 0$.

According to the Macdonald's book [40] $L_{\lambda\mu}$ counts the number of maps f such that $f(\lambda) = \mu$. The following lemma provides an efficient way for computing the matrices $L_{\lambda\mu}$.

Lemma 7.3. *For any λ,μ satisfying $|\lambda| = |\mu|$, denote $v(\lambda) = \ell(\lambda) - m_1(\lambda)$. The following formula holds true:*

$$L_{\lambda\mu} = \sum_{1\leq k_1,\ldots,k_{v(\lambda)}\leq \ell(\mu)} \binom{m_1(\lambda)}{\mu_1 - \sum_{j=1}^{v(\lambda)}\delta_{1,k_j}\lambda_j, \ldots, \mu_{\ell(\mu)} - \sum_{j=1}^{v(\lambda)}\delta_{\ell(\mu),k_j}\lambda_j}. \tag{7.10}$$

Proof. By noticing that the combinatorial meaning of r.h.s. of (7.10) is the same as that of $L_{\lambda\mu}$. □

Remark 7.4. *Let $\kappa_{\lambda\mu} = \frac{L_{\lambda\mu}}{m(\mu)!}$. For any fixed weight $|\lambda| = |\mu|$, $(\kappa_{\lambda\mu})$ is a lower triangular matrix with integer entries which we call* κ-matrix. *Note that we have used the reverse lexicographic ordering for partitions. The first several κ-matrices are given by*

$$(\kappa_{\lambda\mu}) = (1), \quad |\lambda| = |\mu| = 1; \qquad (\kappa_{\lambda\mu}) = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad |\lambda| = |\mu| = 2; \tag{7.11}$$

$$(\kappa_{\lambda\mu}) = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 3 & 1 \end{pmatrix}, \quad |\lambda| = |\mu| = 3; \tag{7.12}$$

$$(\kappa_{\lambda\mu}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 2 & 1 & 1 & 0 \\ 1 & 4 & 3 & 6 & 1 \end{pmatrix}, \quad |\lambda| = |\mu| = 4; \tag{7.13}$$

$$(\kappa_{\lambda\mu}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 2 & 0 & 1 & 0 & 0 \\ 1 & 3 & 4 & 3 & 3 & 1 & 0 \\ 1 & 5 & 10 & 10 & 15 & 10 & 1 \end{pmatrix}, \quad |\lambda| = |\mu| = 5. \tag{7.14}$$

We have the following observations:

(i) The sum of the last row of $(\kappa_{\lambda\mu})$, $|\lambda| = |\mu|$ *is a Bell number. Indeed taking in*

$$\exp\left[\sum_{k\geq 1} h_k(\mathbf{s}) q_k\right] = \sum_{\lambda,\mu\in\mathbb{Y}} \frac{\kappa_{\lambda\mu}}{m(\lambda)!} s_\lambda q_\mu \tag{7.15}$$

$q_1 = q_2 = \ldots = 1$ *we obtain*

$$\exp\left[\exp\left[\sum_{j\geq 1} s_k\right] - 1\right] = \sum_{\lambda\in\mathbb{Y}} \frac{s_\lambda}{m(\lambda)!} \sum_{\mu\in\mathbb{Y}_{|\lambda|}} \kappa_{\lambda\mu}. \tag{7.16}$$

Then taking $s_2 = s_3 = \ldots = 0$ *we obtain*

$$\exp\left(\exp(s_1) - 1\right) = \sum_{k=0} \frac{s_1^k}{k!} \sum_{\mu\in\mathbb{Y}_k} \kappa_{(1^k)\mu}, \tag{7.17}$$

i.e. $\sum_{\mu\in\mathbb{Y}_k} \kappa_{(1^k)\mu} = B_k$ *which is the* k*-th Bell number.*

(ii) The sum of any row of $(\kappa_{\lambda\mu})$, $|\lambda| = |\mu|$ *is a Bell number. More precisely,*

$$\sum_{\mu\in\mathbb{Y}_{|\lambda|}} \kappa_{\lambda\mu} = B_{\ell(\lambda)}. \tag{7.18}$$

Indeed, due to the combinatorial meaning, for a fixed weight the sum only depends on the number of rows of λ*. Since in (ii) we have obtained the sum for* $\lambda = (1^k)$*, the formula (7.18) follows for any* λ*.*

We are now in a position to prove Thm. 1.14.

Proof of Part I. Note that for any partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \ldots) \in \mathbb{Y}$, we have

$$\langle \kappa_{\lambda_1} \ldots \kappa_{\lambda_{\ell(\lambda)}} \tau_{k_1} \ldots \tau_{k_n} \rangle = \frac{\partial^{n+\ell(\lambda)} \log Z^\kappa}{\partial s_{\lambda_1} \ldots \partial s_{\lambda_{\ell(\lambda)}} \partial t_{k_1} \ldots \partial t_{k_n}} (\mathbf{0}; \mathbf{0}). \tag{7.19}$$

According to Theorem 7.1,

$$Z^\kappa(\mathbf{t}; \mathbf{s}) = \mathrm{e}^{-\sum_{k\geq 1} h_k(-\mathbf{s})\, \partial_{t_{k+1}}} Z(\mathbf{t}). \tag{7.20}$$

Therefore we can write the following expansion

$$\log Z^{\kappa}(\mathbf{t};\mathbf{s}) = \sum_{\lambda\in\mathbb{Y}} \frac{(-1)^{\ell(\lambda)} s_\lambda}{m(\lambda)!} \sum_{|\mu|=|\lambda|} L_{\lambda\mu} \frac{(-1)^{\ell(\mu)}}{m(\mu)!} \partial_{t_{\mu_1+1}} \dots \partial_{t_{\mu_{\ell(\mu)}+1}} \log Z(\mathbf{t}). \tag{7.21}$$

From (7.21) we collect the coefficient of s_λ in the generating function $F_n^{\kappa}(z_1,\dots,z_n;\mathbf{s})$. For this purpose we can use the generating formulæ in Thm. 1.2 and Thm. 1.7 to arrive at

$$\begin{aligned}
&\sum_{k_1,\dots,k_n\geq 0} \langle \kappa_{\lambda_1}\dots\kappa_{\lambda_{\ell(\lambda)}}\tau_{k_1}\dots\tau_{k_n}\rangle \frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}} \\
&= (-1)^{\ell(\lambda)} \sum_{|\mu|=|\lambda|} \frac{L_{\lambda\mu}}{m(\mu)!} \operatorname*{res}_{w_1=\infty}\cdots\operatorname*{res}_{w_{\ell(\mu)}=\infty} w^{\mu}\, F_{\ell(\mu)+n}(w_1,\dots,w_{\ell(\mu)},z_1,\dots,z_n)\, dw_1\cdots dw_{\ell(\mu)}
\end{aligned} \tag{7.22}$$

where for a partition μ, $w^{\mu} := \frac{w_1^{2\mu_1+3}}{(2\mu_1+3)!!}\cdots\frac{w_{\ell(\mu)}^{2\mu_{\ell(\mu)}+3}}{(2\mu_{\ell(\mu)}+3)!!}$. This concludes the proof of Part I.

Remark 7.5. *Note that*

$$\partial_{t_{k_1}}\dots\partial_{t_{k_n}} \log Z^{\kappa}(\mathbf{t};\mathbf{s}) = \mathrm{e}^{-\sum_{k\geq 1} h_k(-\mathbf{s})\,\partial_{t_{k+1}}} \partial_{t_{k_1}}\dots\partial_{t_{k_n}} \log Z(\mathbf{t}). \tag{7.23}$$

Let $F_n^{WK}(z_1,\dots,z_n;\mathbf{t})$ and $F_n^{\kappa}(z_1,\dots,z_n;\mathbf{t};\mathbf{s})$ denote the generating functions of n-point correlation functions corresponding to $Z(\mathbf{t})$ and $Z^{\kappa}(\mathbf{t};\mathbf{s})$, respectively. Then we have

$$\begin{aligned}
F_n^{\kappa}(z_1,\dots,z_n;\mathbf{t};\mathbf{s}) &= \mathrm{e}^{-\sum_{k\geq 1} h_k(-\mathbf{s})\,\partial_{t_{k+1}}} F_n^{WK}(z_1,\dots,z_n;\mathbf{t}) \\
&= \sum_{\lambda\in\mathbb{Y}} \frac{(-1)^{\ell(\lambda)} s_\lambda}{m(\lambda)!} \sum_{|\mu|=|\lambda|} L_{\lambda\mu} \frac{(-1)^{\ell(\mu)}}{m(\mu)!} \partial_{t_{\mu_1+1}}\dots\partial_{t_{\mu_{\ell(\mu)}+1}} F_n^{WK}(z_1,\dots,z_n;\mathbf{t}).
\end{aligned} \tag{7.24}$$

Proof of Part II. By definition of the wave function we have

$$\begin{aligned}
\psi^{\kappa}(z;\mathbf{t};\mathbf{s}) &= \frac{Z^{\kappa}(\mathbf{t}-[z^{-1}];\mathbf{s})}{Z^{\kappa}(\mathbf{t};\mathbf{s})} \exp\left(\vartheta(z;\mathbf{t})\right) \\
&= \frac{Z^{\kappa}(\mathbf{t}-[z^{-1}];\mathbf{s})}{Z^{\kappa}(\mathbf{t};\mathbf{s})} \exp\left(\vartheta(z;\mathbf{t})\right) \exp\left(-\sum_{k=2}^{\infty}\frac{h_{k-1}(-\mathbf{s})z^{2k+1}}{(2k+1)!!}\right) \exp\left(\sum_{k=2}^{\infty}\frac{h_{k-1}(-\mathbf{s})z^{2k+1}}{(2k+1)!!}\right) \\
&= \exp\left(\sum_{k=2}^{\infty}\frac{h_{k-1}(-\mathbf{s})z^{2k+1}}{(2k+1)!!}\right) \exp\left(-\sum_{k\geq 1} h_k(-\mathbf{s})\,\partial_{t_{k+1}}\right) \psi^{WK}(z;\mathbf{t}) \\
&= \exp\left(\sum_{k=1}^{\infty}\frac{h_k(-\mathbf{s})z^{2k+3}}{(2k+3)!!}\right) \sum_{\lambda\in\mathbb{Y}} \frac{(-1)^{\ell(\lambda)} s_\lambda}{m(\lambda)!} \sum_{|\mu|=|\lambda|} L_{\lambda\mu} \frac{(-1)^{\ell(\mu)}}{m(\mu)!} \partial_{t_{\mu_1+1}}\dots\partial_{t_{\mu_{\ell(\mu)}+1}} \psi^{WK}(z;\mathbf{t}).
\end{aligned} \tag{7.25}$$

Similarly we have

$$\psi_x^{\kappa}(z;\mathbf{t};\mathbf{s}) = \exp\left(\sum_{k=1}^{\infty}\frac{h_k(-\mathbf{s})z^{2k+3}}{(2k+3)!!}\right) \sum_{\lambda\in\mathbb{Y}} \frac{(-1)^{\ell(\lambda)} s_\lambda}{m(\lambda)!} \sum_{|\mu|=|\lambda|} L_{\lambda\mu} \frac{(-1)^{\ell(\mu)}}{m(\mu)!} \partial_{t_{\mu_1+1}}\dots\partial_{t_{\mu_{\ell(\mu)}+1}} \psi_x^{WK}(z;\mathbf{t}). \tag{7.26}$$

Let

$$A(z;\mathbf{s}) := \psi^{\kappa}(z;\mathbf{0};\mathbf{s}), \quad B(z;\mathbf{s}) := \psi_x^{\kappa}(z;\mathbf{0};\mathbf{s}). \tag{7.27}$$

Note that $Z^\kappa(\mathbf{t};\mathbf{s}) = 1 + \dots$ and that $Z^\kappa(\mathbf{t};\mathbf{0}) = Z(\mathbf{t})$. Expanding in s_1, s_2, $\dots$ we have

$$A(z;\mathbf{s}) = \sum_{k=0}^{\infty} \sum_{\lambda\in\mathbb{Y}_k} A^\lambda(z)\, s_\lambda, \quad A^{(0)}(z) = A^{WK}(z), \tag{7.28}$$

$$B(z;\mathbf{s}) = \sum_{k=0}^{\infty} \sum_{\lambda\in\mathbb{Y}_k} B^\lambda(z)\, s_\lambda, \quad B^{(0)}(z) = B^{WK}(z) \tag{7.29}$$

for some functions $A^\lambda(z)$ and $B^\lambda(z)$. The formulæ (1.48), (1.49) are then obtained from Thm. 1.2, Thm. 1.7 and Thm. 7.1, and this proves Part II. The proof of Thm. 1.14 is complete. □

One can use the formulæ in Lemma 4.1 and the formula (4.20) recursively to obtain

$$\nabla(z_1)\dots\nabla(z_n)\Psi. \tag{7.30}$$

At present we do not have a closed form for these generating functions of "multi-point wave functions".

7.2 Examples of computations of higher Weil–Petersson volumes

In this section we give some explicit examples of calculations of higher Weil–Petersson volumes by using the main theorem and Thm. 1.14.

Example 7.6. (Weil–Petersson volumes). *Consider the special case* $\mathbf{s} = (s, 0, 0, \dots)$. *Then*

$$\log Z^\kappa(x,\mathbf{0};s,\mathbf{0}) = \sum_{g=0}^{\infty} \sum_{\substack{n=0 \\ 3g-3+n\geq 0}}^{\infty} \frac{s^{3g-3+n}}{(3g-3+n)!\,n!} \langle \kappa_1^{3g-3+n}\tau_0^n\rangle_{g,n}\, x^n, \tag{7.31}$$

and by taking twice the x-derivatives we find

$$u_0(x;s) := \partial_x^2 \log Z^\kappa(x,\mathbf{0};s,\mathbf{0}) = \sum_{n\geq 0} \frac{x^n}{n!} \sum_{\substack{g=0 \\ 3g-1+n\geq 0}}^{\infty} \frac{s^{3g-1+n}}{(3g-1+n)!} \langle \kappa_1^{3g-1+n}\tau_0^{n+2}\rangle_{g,n+2}. \tag{7.32}$$

The corresponding n-point functions take the following form

$$F_n^{WP}(z_1,\dots,z_n;s) := \nabla(z_1)\cdots\nabla(z_n)\log Z^\kappa(\mathbf{t};s,\mathbf{0})\,|_{\mathbf{t}=0} = \sum_{g=0}^{\infty} W_{g,n}^{WP}(z_1,\dots,z_n;s) \tag{7.33}$$

where $W_{g,n}^{WP}(z_1,\dots,z_n;s)$ are rational functions of the "Eynard–Orantin type":

$$W_{g,n}^{WP}(z_1,\dots,z_n;s) = \sum_{d=0}^{\infty} \sum_{d+k_1+\dots+k_n=3g-3+n} \langle \kappa_1^d \tau_{k_1}\dots\tau_{k_n}\rangle_g \frac{s^d}{d!} \prod_{i=1}^{n} \frac{(2k_i+1)!!}{z^{2k_i+2}}. \tag{7.34}$$

Define

$$b_{n,g} = \begin{cases} \dfrac{s^{3g-1+n}}{(3g-1+n)!} \langle \kappa_1^{3g-1+n}\tau_0^{n+2}\rangle_{g,n+2}, & \text{if } 3g-1+n\geq 0, \\ 0, & \text{otherwise.} \end{cases} \tag{7.35}$$

An algorithm for computing $u_0(x;s)$ has been given in [57], from a table of which we get

$$b_{0,0}=0,\quad b_{0,1}=\frac{s^2}{8\cdot 2!},\quad b_{0,2}=\frac{787s^5}{128\cdot 5!},\quad \dots, \tag{7.36}$$

$$b_{1,0}=1,\quad b_{1,1}=\frac{7s^3}{6\cdot 3!},\quad b_{1,2}=\frac{1498069s^6}{5760\cdot 6!},\quad \dots, \tag{7.37}$$

$$b_{2,0}=s,\quad b_{2,1}=\frac{529s^4}{24\cdot 4!},\quad b_{2,2}=\frac{10098059s^7}{640\cdot 7!},\quad \dots, \tag{7.38}$$

$$b_{3,0}=5\frac{s^2}{2!},\quad b_{3,1}=\frac{16751s^5}{24\cdot 5!},\quad b_{3,2}=\frac{7473953867s^8}{5760\cdot 8!},\quad \dots. \tag{7.39}$$

Substituting the above initial data into (2.10),(1.28) we obtain

$$W_{0,2}^{WP}=0,\quad W_{1,2}^{WP}=\frac{s^2}{16z_1^2z_2^2}+s\left(\frac{1}{4z_1^4z_2^2}+\frac{1}{4z_2^4z_1^2}\right)+\frac{3}{8z_1^4z_2^4}+\frac{5}{8z_1^6z_2^2}+\frac{5}{8z_1^2z_2^6}, \tag{7.40}$$

$$\begin{aligned}W_{2,2}^{WP}=&\frac{787s^5}{15000z_1^2z_2^2}+s^4\left(\frac{1085}{4608z_1^4z_2^2}+\frac{1085}{4608z_1^2z_2^4}\right)+s^3\left(\frac{551}{576z_1^6z_2^2}+\frac{7}{8z_1^4z_2^4}+\frac{551}{576z_1^2z_2^6}\right)\\&+s^2\left(\frac{399}{128z_1^8z_2^2}+\frac{181}{64z_1^6z_2^4}+\frac{181}{64z_1^4z_2^6}+\frac{399}{128z_1^2z_2^8}\right)\\&+s\left(\frac{231}{32z_1^{10}z_2^2}+\frac{203}{32z_1^8z_2^4}+\frac{105}{16z_1^6z_2^6}+\frac{203}{32z_1^4z_2^8}+\frac{231}{32z_1^2z_2^{10}}\right)+\frac{1155}{128z_1^{12}z_2^2}+\frac{945}{128z_1^{10}z_2^4}\\&+\frac{1015}{128z_1^8z_2^6}+\frac{1015}{128z_1^6z_2^8}+\frac{945}{128z_1^4z_2^{10}}+\frac{1155}{128z_1^2z_2^{12}},\end{aligned} \tag{7.41}$$

$$W_{0,3}^{WP}=\frac{1}{z_1^2z_2^2z_3^2}, \tag{7.42}$$

$$\begin{aligned}W_{1,3}^{WP}=&\frac{7s^3}{36z_1^2z_2^2z_3^2}+s^2\left(\frac{13}{16z_1^4z_2^2z_3^2}+\frac{13}{16z_1^2z_2^4z_3^2}+\frac{13}{16z_1^2z_2^2z_3^4}\right)\\&+s\left(\frac{5}{2z_1^6z_2^2z_3^2}+\frac{9}{4z_1^4z_2^4z_3^2}+\frac{9}{4z_1^4z_2^2z_3^4}+\frac{5}{2z_1^2z_2^6z_3^2}+\frac{9}{4z_1^2z_2^4z_3^4}+\frac{5}{2z_1^2z_2^2z_3^6}\right)\\&+\frac{35}{8z_1^8z_2^2z_3^2}+\frac{15}{4z_1^6z_2^4z_3^2}+\frac{15}{4z_1^6z_2^2z_3^4}+\frac{15}{4z_1^4z_2^6z_3^2}+\frac{9}{4z_1^4z_2^4z_3^4}+\frac{15}{4z_1^4z_2^2z_3^6}\\&+\frac{35}{8z_1^2z_2^8z_3^2}+\frac{15}{4z_1^2z_2^6z_3^4}+\frac{15}{4z_1^2z_2^4z_3^6}+\frac{35}{8z_1^2z_2^2z_3^8}.\end{aligned} \tag{7.43}$$

The expressions for $W_{1,2}^{WP}, W_{0,3}^{WP}$ coincide with those derived in [54]; the function $W_{0,2}^{WP}$ in [54] is not vanishing since some natural supplementary definitions for intersection numbers are used in [54]. Let $Vol_{g,n}(L_1,\dots,L_n):=Vol(\mathcal{M}_{g,n}(L_1,\dots,L_n))$ denote the Weil–Petersson volumes and let

$$v_{g,n}(L_1,\dots,L_n):=\frac{Vol_{g,n}(L_1,\dots,L_n)}{(2\pi)^{3g-3+n}}. \tag{7.44}$$

It was shown in [43, 44] that

$$v_{g,n}(L_1,\dots,L_n)=\sum_{d+k_1+\dots k_n=3g-3+n}\frac{\langle\kappa_1^d\tau_{k_1}\dots\tau_{k_n}\rangle_g}{d!\,k_1!\cdots k_n!}\,L_1^{2k_1}\cdots L_n^{2k_n}. \tag{7.45}$$

We remark that the relationship between $v_{g,n}$ and $W_{g,n}^{WP}$ is a Laplace transform [46, 24, 53]:

$$W_{g,n}^{WP}(z_1,\dots,z_n;s=1)=2^n\int_0^\infty\cdots\int_0^\infty L_1\cdots L_n\cdot v_{g,n}(L_1,\dots,L_n)\cdot \mathrm{e}^{-\sum_{i=1}^n\sqrt{2}z_iL_i}\,dL_1\dots dL_n. \tag{7.46}$$

Note that in the above example we have not used Thm. 1.14. More generally, given any initial value problem of the KdV hierarchy with an initial data $u(x,\mathbf{0}) = u_0(x) \in \mathbb{C}[[x]]$ one can compute the resolvent function $\mathcal{R}(z;x)$ of the Lax operator by solving the ODE (2.11). Then, with the help of the Main Theorem, the generating functions of multipoint correlators readily follow.

Let us now give examples of application of Thm. 1.14.

Example 7.7. (linear insertion of κ-classes). *For $n = 0$, $\lambda = (j)$ we have*

$$\langle \kappa_j \rangle = \operatorname*{res}_{w=\infty} \frac{-w^{2j+3}}{(2j+3)!!} F_1(w)\, dw = \operatorname*{res}_{w=\infty} \frac{-w^{2j+3}}{(2j+3)!!} \sum_{g=1}^{\infty} \frac{(6g-3)!!}{24^g g!} w^{-(6g-2)} dw. \tag{7.47}$$

So we have

$$\langle \kappa_{3g-3} \rangle_{g,0} = \frac{1}{24^g \cdot g!}. \tag{7.48}$$

For $n = 1$, $\lambda = (j)$ we have

$$\sum_{k_1 \geq 0} \langle \kappa_j \tau_{k_1} \rangle \frac{(2k_1+1)!!}{z^{2k_1+2}} = \operatorname*{res}_{w=\infty} \frac{-w^{2j+3}}{(2j+3)!!} F_2(w,z)\, dw \tag{7.49}$$

$$= \operatorname*{res}_{w=\infty} \frac{-w^{2j+3}}{(2j+3)!!} \left(\frac{\operatorname{Tr}\left(M(w)M(z)\right)}{(w^2-z^2)^2} - \frac{w^2+z^2}{(w^2-z^2)^2} \right) dw \tag{7.50}$$

$$= \frac{1}{(2j+3)!!} \operatorname{Tr} \left(\frac{1}{2z} \left[\partial_z \left(z^{2j+2} M(z) \right) \right]_+ \cdot M(z) \right) - \frac{1}{(2j+1)!!} z^{2j+2}. \tag{7.51}$$

Here "$+$" means taking the polynomial part in the z expansion at infinity. Particularly, if $n = 1$, $j = 1$ then we have

$$\frac{1}{2z} \left[\partial_z \left(z^4 M(z) \right) \right]_+ = \begin{pmatrix} -\frac{1}{2} & -2z^2 \\ -3z^4 & \frac{1}{2} \end{pmatrix}; \tag{7.52}$$

hence

$$\sum_{k \geq 0} \langle \kappa_1 \tau_k \rangle \frac{(2k+1)!!}{z^{2k+2}} = \frac{1}{5!!} \operatorname{Tr} \left(\frac{1}{2z} \left[\partial_z \left(z^4 M(z) \right) \right]_+ \cdot M(z) \right) - \frac{1}{3!!} z^4$$

$$= \frac{1}{5!!} \sum_{g=1}^{\infty} \left(\frac{1}{2} \frac{(6g-5)!!}{24^{g-1} \cdot (g-1)!} - 2 \frac{6g+1}{6g-1} \frac{(6g-1)!!}{24^g \cdot g!} + 3 \frac{(6g-1)!!}{24^g \cdot g!} \right) z^{-6g+4}$$

$$= 3 \sum_{g=1}^{\infty} \frac{(12g^2-12g+5)(6g-5)!!}{5!! \cdot 24^g \cdot g!} z^{-6g+4}. \tag{7.53}$$

This means that

$$\langle \kappa_1 \tau_{3g-3} \rangle_{g,1} = 3 \frac{12g^2-12g+5}{5!! \cdot 24^g \cdot g!},\ g \geq 1. \tag{7.54}$$

If $n = 1$, $j = 2$ then we have

$$\frac{1}{2z} \left[\partial_z \left(z^6 M(z) \right) \right]_+ = \begin{pmatrix} -z^2 & -3z^4 \\ \frac{7}{8} - 4z^6 & z^2 \end{pmatrix}; \tag{7.55}$$

whence a straightforward manipulation of series shows

$$\sum_{k\geq 0}\langle\kappa_2\tau_k\rangle\frac{(2k+1)!!}{z^{2k+2}}=\frac{1}{7!!}\mathrm{Tr}\left(\frac{1}{2z}\left[\partial_z\left(z^6M(z)\right)\right]_+\cdot M(z)\right)-\frac{1}{5!!}z^6$$
$$=3\sum_{g=1}^{\infty}\frac{(72g^3-132g^2+95g-35)(6g-7)!!}{7!!\cdot 24^g\cdot g!}z^{-6g+6}. \tag{7.56}$$

This means that

$$\langle\kappa_2\tau_{3g-4}\rangle_{g,1}=3\frac{72g^3-132g^2+95g-35}{7!!\cdot 24^g\cdot g!},\qquad g\geq 2. \tag{7.57}$$

Similar computations lead to

$$\langle\kappa_3\tau_{3g-5}\rangle_{g,1}=\frac{1296g^4-3888g^3+4482g^2-2835g+945}{9!!\cdot 24^g\cdot g!},\qquad g\geq 2. \tag{7.58}$$

In general, we have

$$\langle\kappa_j\tau_{3g-j-2}\rangle_{g,1}\sim\frac{6^{j+1}g^{j+1}}{(2j+3)!!\cdot 24^g\cdot g!},\quad g\to\infty. \tag{7.59}$$

For $n\geq 2$, $\lambda=(j)$ we have

$$\sum_{k_1,\ldots,k_n\geq 0}\langle\kappa_j\tau_{k_1}\ldots\tau_{k_n}\rangle\frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}}=\operatorname*{res}_{w=\infty}\frac{-w^{2j+3}}{(2j+3)!!}F_{n+1}(w,z_1,\ldots,z_n)\,dw \tag{7.60}$$
$$=\operatorname*{res}_{w=\infty}\frac{w^{2j+3}}{(2j+3)!!}\sum_{r\in S_n}\frac{\mathrm{Tr}\left(M(w)M(z_{r_1})\cdots M(z_{r_n})\right)}{(w^2-z_{r_1}^2)(z_{r_n}^2-w^2)\prod_{j=1}^{n-1}(z_{r_j}^2-z_{r_{j+1}}^2)} \tag{7.61}$$
$$=\frac{1}{(2j+3)!!}\operatorname*{res}_{w=\infty}\sum_{r\in S_n}\frac{\mathrm{Tr}\left(M(w)M(z_{r_1})\cdots M(z_{r_n})\right)}{\prod_{j=1}^{n}(z_{r_j}^2-z_{r_{j+1}}^2)}\left(\frac{w^{2j+3}}{w^2-z_{r_1}^2}-\frac{w^{2j+3}}{w^2-z_{r_n}^2}\right)dw \tag{7.62}$$
$$=-\frac{1}{(2j+3)!!}\sum_{r\in S_n}\frac{\mathrm{Tr}\left((z_{r_1}^{2j+2}M(z_{r_1}))_+-(z_n^{2j+2}M(z_{r_n}))_+)M(z_{r_1})\cdots M(z_{r_n})\right)}{\prod_{j=1}^{n}(z_{r_j}^2-z_{r_{j+1}}^2)}. \tag{7.63}$$

So we can also collect the following generating function for linear insertion of κ-classes:

$$\sum_{j\geq 1}\sum_{k_1,\ldots,k_n\geq 0}\langle\kappa_j\tau_{k_1}\ldots\tau_{k_n}\rangle\frac{(2j+3)!!}{w^{2j}}\frac{(2k_1+1)!!}{z_1^{2k_1+2}}\cdots\frac{(2k_n+1)!!}{z_n^{2k_n+2}}=\left[w^4F_{n+1}(w,z_1,\ldots,z_n)\right]_- \tag{7.64}$$

where "$-$" means taking the negative part in the w expansion at ∞.

Example 7.8 (linear deformation of the wave function). *For $j\geq 1$, we have*

$$A^{(j)}(z) = -\frac{z^{2j+3}}{(2j+3)!!}A^{(0)}(z)-\sum_{|\mu|=j}L_{(j),\mu}\frac{(-1)^{\ell(\mu)}}{m(\mu)!}\partial_{t_{\mu_1+1}}\ldots\partial_{t_{\mu_{\ell(\mu)}+1}}\psi^{WK}(z;\mathbf{t})|_{\mathbf{t}=\mathbf{0}}$$
$$= -\frac{z^{2j+3}}{(2j+3)!!}c(z)+\psi^{WK}_{t_{j+1}}(z;\mathbf{0}). \tag{7.65}$$

By using Lemma 4.1 and Corollary 6.4 we obtain

$$\begin{aligned}
&\nabla(w)\psi^{WK}(z;\mathbf{t})|_{\mathbf{t}=\mathbf{0}}\\
&=\frac{2S(w;\mathbf{0})B^{WK}(z)-\mathcal{R}_x(w;\mathbf{0})A^{WK}(z)}{2\,(w^2-z^2)}\\
&=\frac{2\sum_{g=0}^{\infty}\frac{(6g-1)!!}{24^g\cdot g!}w^{-6g}z\,q(z)-\sum_{g=1}^{\infty}\frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!}w^{-6g+4}c(z)}{2\,(w^2-z^2)}\\
&=\frac{z\,q}{w^2}+\frac{2z^3q-c}{2\,w^4}+\frac{4z^5q-2z^2c}{4\,w^6}+\frac{(8z^7+5z)\,q-4z^4c}{8\,w^8}+\frac{(16z^9+10z^3)q-(8z^6+35)c}{16\,w^{10}}+\mathcal{O}(w^{-12}).
\end{aligned}\tag{7.66}$$

So we have

$$A^{(1)}(z)=-\frac{z^5}{5!!}c+\frac{z^5}{5!!}q-\frac{z^2}{2\cdot 5!!}c=-\frac{1}{24}z^{-1}+\frac{77}{576}z^{-4}+\mathcal{O}(z^{-7}),\tag{7.67}$$

$$A^{(2)}(z)=-\frac{z^7}{7!!}c+\frac{8z^7+5z}{8\cdot 7!!}q-\frac{z^4}{2\cdot 7!!}c=\frac{1}{48}z^{-2}-\frac{13}{144}z^{-5}+\mathcal{O}(z^{-8}),\tag{7.68}$$

$$A^{(3)}(z)=-\frac{z^9}{9!!}c+\frac{16z^9+10z^3}{16\cdot 9!!}q-\frac{8z^6+35}{16\cdot 9!!}c=-\frac{11}{1152}z^{-3}+\frac{1639}{27648}z^{-6}+\mathcal{O}(z^{-9}),\tag{7.69}$$

etc. In the above formulæ it is understood that $q=q(z)$, $c=c(z)$ are Faber–Zagier series. As it was expected, indeed, all deformation terms $A^{\lambda}(z)$, $\lambda\neq(0)$ contain only negative powers in their expansions at $z\to\infty$. Similarly, we have

$$B^{(j)}(z)=-\frac{z^{2j+4}}{(2j+3)!!}q(z)+\psi^{WK}_{t_0t_{j+1}}(z;\mathbf{0}),\tag{7.70}$$

$$\begin{aligned}
&\nabla(w)\psi^{WK}_x(z;\mathbf{t})|_{\mathbf{t}=\mathbf{0}}=\frac{\mathcal{R}_x(w;\mathbf{0})B^{WK}(z)-\left[\mathcal{R}_{xx}(w;\mathbf{0})-2S(w;\mathbf{0})z^2\right]A^{WK}(z)}{2\,(w^2-z^2)}\\
&=\frac{\sum_{g=1}^{\infty}\frac{(6g-5)!!}{24^{g-1}\cdot(g-1)!}w^{-6g+4}z\,q(z)-\left[\sum_{g=1}^{\infty}\frac{(6g-3)!!}{24^{g-1}\cdot(g-1)!}w^{-6g+2}-2\sum_{g=0}^{\infty}\frac{(6g-1)!!}{24^g\cdot g!}w^{-6g}z^2\right]c(z)}{2\,(w^2-z^2)}\\
&=\frac{z^2c}{w^2}+\frac{2z^4c+zq}{2w^4}+\frac{(4z^6-6)c+2z^3q}{4w^6}+\frac{(8z^8-7z^2)c+4z^5q}{8w^8}+\mathcal{O}(w^{-10}).
\end{aligned}\tag{7.71}$$

And we have

$$B^{(1)}(z)=-\frac{z^6}{5!!}q+\frac{z^3}{2\cdot 5!!}q+\frac{4z^6-6}{4\cdot 5!!}c=-\frac{1}{24}+\frac{79}{576}z^{-3}+\frac{18095}{27648}z^{-6}+\mathcal{O}(z^{-9}),\tag{7.72}$$

$$B^{(2)}(z)=-\frac{z^8}{7!!}q+\frac{z^5}{2\cdot 7!!}q+\frac{8z^8-7z^2}{8\cdot 7!!}c=-\frac{1}{48}z^{-1}-\frac{55}{576}z^{-4}-\frac{31603}{55296}z^{-7}+\mathcal{O}(z^{-10}).\tag{7.73}$$

These expressions agree with expression derived from a less straightforward method in Appendix B.

Example 7.9 (**higher deformation of the wave function**). *Consider deformation of the wave function associated to partitions of form (1^k), $k\geq 1$. For $k=1$, it has been solved above. Let $k=2$; we have*

$$\begin{aligned}
A^{(1^2)}(z)&=\left(\frac{z^7}{2\cdot 7!!}+\frac{z^{10}}{2\cdot 5!!^2}\right)c-\frac{z^5}{5!!}\psi^{WK}_{t_2}(z;\mathbf{0})+\frac{1}{2!}\sum_{|\mu|=2}L_{(1^2),\mu}\frac{(-1)^{\ell(\mu)}}{m(\mu)!}\partial_{t_{\mu_1+1}}\dots\partial_{t_{\mu_{\ell(\mu)}+1}}\psi^{WK}(z;\mathbf{t})|_{\mathbf{t}=\mathbf{0}}\\
&=\left(\frac{z^7}{2\cdot 7!!}+\frac{z^{10}}{2\cdot 5!!^2}\right)c-\frac{z^5}{5!!}\frac{4z^5q-2z^2c}{4\cdot 5!!}+\frac{1}{2!}\left(-\frac{(8z^7+5z)\,q-4z^4c}{8\cdot 7!!}+\psi^{WK}_{t_2t_2}(z;\mathbf{0})\right)
\end{aligned}\tag{7.74}$$

and

$$B^{(1^2)}(z) = \left(\frac{z^8}{2\cdot 7!!} + \frac{z^{11}}{2\cdot 5!!^2}\right) q - \frac{z^5}{5!!}\psi^{WK}_{t_0t_2}(z;\mathbf{0}) + \frac{1}{2!}\sum_{|\mu|=2} L_{(1^2),\mu}\frac{(-1)^{\ell(\mu)}}{m(\mu)!}\partial_{t_{\mu_1+1}}\dots\partial_{t_{\mu_{\ell(\mu)}+1}}\psi^{WK}_x(z;\mathbf{t})|_{\mathbf{t}=\mathbf{0}}$$

$$= \left(\frac{z^8}{2\cdot 7!!} + \frac{z^{11}}{2\cdot 5!!^2}\right) q - \frac{z^5}{5!!}\frac{(4z^6-6)c + 2z^3 q}{4\cdot 5!!} + \frac{1}{2!}\left(-\frac{(8z^8-7z^2)c + 4z^5 q}{8\cdot 7!!} + \psi^{WK}_{t_0t_2t_2}(z;\mathbf{0})\right). \tag{7.75}$$

By using (4.1) one can derive that

$$\psi^{WK}_{t_2t_2}(z;\mathbf{0}) = \frac{4z^{10}-5z^4}{4\cdot 5!!^2}c + \frac{z}{4\cdot 5!!}q, \quad \psi^{WK}_{t_0t_2t_2}(z;\mathbf{0}) = \frac{z^{11}q}{5!!^2} - \frac{z^5 q}{180} - \frac{z^2 c}{60}. \tag{7.76}$$

So we have

$$\begin{aligned} A^{(1^2)}(z) &= \left(\frac{z^{10}}{225} + \frac{11z^7}{1575} - \frac{z^4}{2520}\right)c + \left(-\frac{z^{10}}{225} - \frac{z^7}{210} + \frac{3z}{560}\right)q \\ &= \frac{37}{1152}z^{-2} - \frac{28249}{138240}z^{-5} + \mathcal{O}(z^{-8}), \end{aligned} \tag{7.77}$$

$$\begin{aligned} B^{(1^2)}(z) &= \left(-\frac{z^{11}}{225} - \frac{z^8}{210} + \frac{z^5}{150} - \frac{z^2}{240}\right)c + \left(\frac{z^{11}}{225} + \frac{4z^8}{1575} - \frac{13z^5}{2520}\right)q \\ &= -\frac{35}{1152}z^{-1} + \frac{29051}{138240}z^{-4} + \mathcal{O}(z^{-7}). \end{aligned} \tag{7.78}$$

Using Part I of Theorem 1.14 we find

$$\langle\kappa_1^2\tau_2\rangle_{2,1} = \frac{139}{11520}, \quad \langle\kappa_1^2\tau_5\rangle_{3,1} = \frac{3781}{2903040}, \quad \langle\kappa_1^2\tau_8\rangle_{4,1} = \frac{48689}{928972800} \tag{7.79}$$

and

$$\begin{aligned} F_2^\kappa(z,w;s_1=s,\mathbf{0}) &= s^2\left(\frac{300825}{1024w^8z^8} + \frac{399}{128w^8z^2} + \frac{181}{64w^6z^4} + \frac{181}{64w^4z^6} + \frac{399}{128w^2z^8} + \frac{1}{16w^2z^2} + \dots\right) \\ &\quad + s\left(\frac{231}{32w^{10}z^2} + \frac{203}{32w^8z^4} + \frac{105}{16w^6z^6} + \frac{203}{32w^4z^8} + \frac{1}{4w^4z^2} + \frac{231}{32w^2z^{10}} + \frac{1}{4w^2z^4} + \dots\right) \\ &\quad + \left(\frac{1015}{128w^8z^6} + \frac{1015}{128w^6z^8} + \frac{5}{8w^6z^2} + \frac{3}{8w^4z^4} + \frac{5}{8w^2z^6} + \dots\right) + o(s^2). \end{aligned} \tag{7.80}$$

An alternative method for computing $A(z;\mathbf{s}), B(z;\mathbf{s})$ is presented in Appendix B.

8 Further remarks

It would be interesting to prove directly the equivalence between the formula (1.36), the (explicit) integral/recursive formulæ of "n-point functions" given by Okounkov [48], Liu–Xu [36, 37], Brézin–Hikami [10, 11], and Kontsevich's main identity [34]. We indicate the relation between these three as follows. Recall that, given a formal (divergent) series of the form

$$f(z) = \sum_{k=0}^{\infty} v_k \frac{(2k+1)!!}{z^{2k+2}}, \tag{8.1}$$

it can be re-summed by a suitable version of the Borel summation method. Noting that

$$(2k+1)!! = \frac{2^{k+1}}{\sqrt{\pi}} \int_0^\infty u^{\frac{2k+1}{2}} \mathrm{e}^{-u} du, \quad k \geq 0 \tag{8.2}$$

we have, integrating term-by-term

$$\sum_{k=0}^{\infty} v_k \frac{(2k+1)!!}{z^{2k+2}} = \frac{2\,z}{\sqrt{\pi}} \int_0^\infty s^{\frac{1}{2}}\, \hat{f}(s)\, \mathrm{e}^{-sz^2}\, ds \tag{8.3}$$

where $\hat{f}(s) := \sum_{k=0}^{\infty} v_k (2s)^k$ is the Borel re-summation of $f(z)$. In general we say that $f(z)$ is Borel summable if the function $\hat{f}(s)$ (the "Borel transform" of f) has a nonzero radius of convergence, and it can be extended analytically along a strip surrounding the real positive axis. It then follows that the relation between the generating functions of intersection numbers that we have constructed in this paper and Okounkov's generating functions is precisely that the latter are Borel transforms of the former, or, which is the same, that the former are asymptotic expansions of the Laplace transforms of the latter. The relation between our generating function and Kontsevich's is simpler. To be precise, let us introduce the following notations

$$F_n^{OK}(x_1, \dots, x_n) = \sum_{k_1,\dots,k_n=0}^{\infty} \langle \tau_{k_1} \dots \tau_{k_n} \rangle\, x_1^{k_1} \cdots x_n^{k_n}, \tag{8.4}$$

$$F_n^{K}(z_1, \dots, z_n) = \sum_{k_1,\dots,k_n=0}^{\infty} \frac{(2k_1-1)!!}{z_1^{2k_1+2}} \cdots \frac{(2k_n-1)!!}{z_n^{2k_n+2}} \langle \tau_{k_1} \dots \tau_{k_n} \rangle. \tag{8.5}$$

Then we have

$$F_n^{WK}(z_1, \dots, z_n) = \frac{2^n\, z_1 \cdots z_n}{\pi^{\frac{n}{2}}} \int_0^\infty \cdots \int_0^\infty (x_1 \cdots x_n)^{\frac{1}{2}}\, F_n^{OK}(2x_1, \dots, 2x_n)\, \mathrm{e}^{-x_1 z_1^2 - \dots - x_n z_n^2}\, dx_1 \cdots dx_n, \tag{8.6}$$

$$F_n^{WK}(z_1, \dots, z_n) = (-1)^n \left(\prod_{j=1}^{n} \frac{\partial}{\partial z_j} z_j \right) \left(F_n^K(z_1, \dots, z_n) \right). \tag{8.7}$$

For $n = 1$, one can directly verify (8.6) (actually in the case $n = 1$ the generating series of Okounkov and ours are both well-known). Indeed,

$$F_1^{OK}(2x_1) = \frac{1}{4x_1^2}\left(\mathrm{e}^{\frac{x_1^3}{3}} - 1\right) = \sum_{g=1}^{\infty} \frac{1}{24^g g!} (2x_1)^{3g-2}. \tag{8.8}$$

Recall that

$$F_1^{WK}(z_1) = \sum_{g=1}^{\infty} \frac{(6g-3)!!}{24^g g!} z_1^{-(6g-2)}. \tag{8.9}$$

So it is straightforward to verify (8.6) for $n = 1$. However, it appears that for $n \geq 2$ a direct verification of (8.6) is not trivial.

The final remark is that the formula (1.28) for multi-point correlation functions possesses certain universality in tau-symmetric integrable systems. Indeed, it can be generalized to the Gelfand-Dickey hierarchy and more generally to the Drinfeld-Sokolov hierarchy, as well as to the Jimbo–Miwa–Ueno isomonodromic problems [7, 29], which will be presented in a separate publication [8]. It would be interesting to investigate whether the formula works for all integrable hierarchies of topological (or cohomological) type associated to semisimple Frobenius manifolds [22, 21]: the first nontrivial examples in this investigation would be the intermediate long wave hierarchy [13], the discrete KdV hierarchy [21] and the extended Toda hierarchy [14].

A Some useful formulæ

The wave functions associated to the Witten–Kontsevich tau-function $Z(\mathbf{t})$ at $t_1 = t_2 = \ldots = 0$ satisfy

$$\psi(z; t_0, 0, 0, \ldots) = \sqrt{2\pi z}\, \mathrm{e}^{\frac{z^3}{3}} 2^{\frac{1}{3}}\, \mathrm{Ai}\,(\xi)\,, \tag{A.1}$$

$$\psi^*(z; t_0, 0, 0, \ldots) = \mathrm{e}^{\frac{\pi i}{6}} \sqrt{2\pi z}\, \mathrm{e}^{-\frac{z^3}{3}} 2^{\frac{1}{3}}\, \mathrm{Ai}\,(\omega\xi)\,, \tag{A.2}$$

where $\xi = 2^{-\frac{2}{3}} \left(z^2 - 2\, t_0\right),\ \omega = \mathrm{e}^{\frac{2\pi\sqrt{-1}}{3}}$. Noting that

$$\xi_x = -2^{\frac{1}{3}}, \quad \xi_z = 2^{\frac{1}{3}} z \tag{A.3}$$

we have

$$\psi_x(z; t_0, 0, 0, \ldots) = -\sqrt{2\pi z}\, \mathrm{e}^{\frac{z^3}{3}} 2^{\frac{2}{3}}\, \mathrm{Ai}'(\xi), \tag{A.4}$$

$$\psi^*_x(z; t_0, 0, 0, \ldots) = -\mathrm{e}^{\frac{5}{6}\pi\sqrt{-1}} \sqrt{2\pi z}\, \mathrm{e}^{-\frac{z^3}{3}} 2^{\frac{2}{3}}\, \mathrm{Ai}'(\omega\xi), \tag{A.5}$$

$$\psi_z(z; t_0, 0, 0, \ldots) = \sqrt{2\pi z}\, \mathrm{e}^{\frac{z^3}{3}} 2^{\frac{1}{3}} \left[\left(\frac{1}{2z} + z^2\right) \mathrm{Ai}(\xi) + 2^{\frac{1}{3}} z\, \mathrm{Ai}'(\xi)\right], \tag{A.6}$$

$$\psi^*_z(z; t_0, 0, 0, \ldots) = -\sqrt{2\pi z}\, \mathrm{e}^{\frac{-z^3}{3}} 2^{\frac{1}{3}} \left[\mathrm{e}^{\frac{1}{6}\pi i} \left(\frac{1}{-2z} + z^2\right) \mathrm{Ai}(\omega\xi) - \mathrm{e}^{\frac{5}{6}\pi i} 2^{\frac{1}{3}} z\, \mathrm{Ai}'(\omega\xi)\right], \tag{A.7}$$

$$\psi_{zx}(z; t_0, 0, 0, \ldots) = -\sqrt{2\pi z}\, \mathrm{e}^{\frac{z^3}{3}} 2^{\frac{2}{3}} \left[\left(\frac{1}{2z} + z^2\right) \mathrm{Ai}'(\xi) + 2^{-\frac{1}{3}} z \left(z^2 - 2t_0\right) \mathrm{Ai}(\xi)\right], \tag{A.8}$$

$$\psi^*_{zx}(z; t_0, 0, 0, \ldots) = \sqrt{2\pi z}\, \mathrm{e}^{\frac{-z^3}{3}} 2^{\frac{2}{3}} \left[\mathrm{e}^{\frac{5}{6}\pi i} \left(\frac{1}{-2z} + z^2\right) \mathrm{Ai}'(\omega\xi) - \mathrm{e}^{\frac{1}{6}\pi i} 2^{-\frac{1}{3}} z \left(z^2 - 2t_0\right) \mathrm{Ai}(\omega\xi)\right]. \tag{A.9}$$

B Generalized Kac–Schwarz operator and higher Weil–Petersson volumes

Let us first compute $A(z; \mathbf{s}) = \psi^\kappa(z; \mathbf{0}; \mathbf{s})$, $B(z; \mathbf{s}) = \psi^\kappa_x(z; \mathbf{0}; \mathbf{s})$ associated with higher Weil–Petersson volumes by using the technique of generalized Kac–Schwarz operators explained above.

Theorem 7.1 together with eq. (6.3) implies that $Z^\kappa(\mathbf{t}; \mathbf{s})$ satisfies the string equation

$$\sum_{k\geq 0} \tilde{t}_{k+1} \frac{\partial Z^\kappa}{\partial t_k} + \frac{\tilde{t}_0^2}{2} Z^\kappa = 0 \tag{B.1}$$

where $\tilde{t}_0 = t_0$, $\tilde{t}_k = t_k - h_{k-1}(-\mathbf{s})$, $k \geq 1$, namely,

$$c_0 = 0, \quad c_k = h_{k-1}(-\mathbf{s}),\ k \geq 1. \tag{B.2}$$

The corresponding generalized Kac–Schwarz operator (see Definition 5.1) reads as follows

$$S_z = \frac{1}{z}\partial_z - \frac{1}{2z^2} - z - \sum_{k=1}^{\infty} \frac{h_k(-\mathbf{s})}{(2k+1)!!} z^{2k+1} = S_z^{WK} - \sum_{k=1}^{\infty} \frac{h_k(-\mathbf{s})}{(2k+1)!!} z^{2k+1}. \tag{B.3}$$

As before let $f(z; x; \mathbf{s}) = \psi^\kappa(z; x, \mathbf{0}; \mathbf{s})$, $u_0(x; \mathbf{s}) = \partial_x^2 \log Z^\kappa(x, \mathbf{0}; \mathbf{s})$. Then according to Prop. 5.3 we have

$$S_z f(z; x; \mathbf{s}) = -f_x(z; x; \mathbf{s}) - \sum_{k\geq 1} h_k(-\mathbf{s}) \psi^\kappa_{t_k}(z; x, \mathbf{0}; \mathbf{s}), \tag{B.4}$$

$$S_z f_x(z;x;\mathbf{s}) = -(z^2 - 2\,u_0(x;\mathbf{s}))f(z;x;\mathbf{s}) - \sum_{k\geq 1} h_k(-\mathbf{s})\psi^{\kappa}_{t_0t_k}(z;x,\mathbf{0};\mathbf{s}), \tag{B.5}$$

$$\psi^{\kappa}_{t_k}(z;x,\mathbf{0};\mathbf{s}) = \frac{1}{(2k+1)!!}\sum_{i=0}^{k}(2i-1)!!\, z^{2k-2i}\left(\Omega_{i-1;0}|_{u\to u_0} f_x(z;x;\mathbf{s}) - \frac{1}{2}f(z;x;\mathbf{s})\partial_x\Omega_{i-1;0}|_{u\to u_0}\right), \tag{B.6}$$

$$\begin{aligned}\psi^{\kappa}_{t_0t_k}(z;x,\mathbf{0}) = \frac{1}{(2k+1)!!}\sum_{i=0}^{k}(2i-1)!!\, z^{2k-2i}&\left(\partial_x\Omega_{i-1;0}|_{u\to u_0} f_x(z;x) + \Omega_{i-1;0}|_{u\to u_0}\left(z^2 - 2u_0(x)\right) f(z;x)\right.\\ &\left. -\frac{1}{2}f_x(z;x)\partial_x\Omega_{i-1;0}|_{u\to u_0} - \frac{1}{2}f(z;x)\partial_x^2\Omega_{i-1;0}|_{u\to u_0}\right)\end{aligned} \tag{B.7}$$

Taking $x = 0$ in the above equations (B.4),(B.5) we arrive at

Lemma B.1. *The functions $A(z;\mathbf{s})$ and $B(z;\mathbf{s})$ satisfy*

$$S_z A(z;\mathbf{s}) = -B(z;\mathbf{s}) - \sum_{k\geq 1} h_k(-\mathbf{s})\,\psi^{\kappa}_{t_k}(z;\mathbf{0};\mathbf{s}), \tag{B.8}$$

$$S_z B(z;\mathbf{s}) = -(z^2 - 2\,u_0(0;\mathbf{s}))A(z;\mathbf{s}) - \sum_{k\geq 1} h_k(-\mathbf{s})\,\psi^{\kappa}_{t_0t_k}(z;\mathbf{0};\mathbf{s}). \tag{B.9}$$

To solve the above equations (B.8), (B.9) we can expand them as formal series in $\mathbf{s}$ and compare the coefficients. At the linear approximation in $\mathbf{s}$, employing Lemma 6.2 and comparing the coefficients of s_j, $j \geq 1$, we get

$$\begin{aligned}\frac{z^{2j+1}}{(2j+1)!!}A^{(0)}(z) + S_z^{WK}A^{(j)}(z) &= -B^{(j)}(z) + \frac{z^{2j}}{(2j+1)!!}B^{(0)}(z)\\ &\quad + \sum_{i=1}^{j}\frac{(2i-1)!!}{(2j+1)!!}z^{2j-2i}\left(\langle\tau_{i-1}\tau_0\rangle\, B^{(0)}(z) - \frac{1}{2}\langle\tau_{i-1}\tau_0^2\rangle A^{(0)}(z)\right),\end{aligned} \tag{B.10}$$

$$\begin{aligned}\frac{z^{2j+1}}{(2j+1)!!}B^{(0)}(z) + S_z^{WK}B^{(j)}(z) &= -z^2A^{(j)}(z) + 2\langle\tau_0^2\kappa_j\rangle A^{(j)}(z) + \frac{z^{2j+2}}{(2j+1)!!}A^{(0)}(z)\\ &\quad + \sum_{i=1}^{j}\frac{(2i-1)!!}{(2j+1)!!}z^{2j-2i}\left(\frac{1}{2}\langle\tau_0^2\tau_{i-1}\rangle B^{(0)}(z) + \langle\tau_0\tau_{i-1}\rangle z^2A^{(0)}(z) - \frac{1}{2}A^{(0)}(z)\langle\tau_0^3\tau_{i-1}\rangle\right).\end{aligned} \tag{B.11}$$

Also, taking derivatives w.r.t. x in the string equation (B.1) and taking $\mathbf{t} = 0$ yields

Lemma B.2. *For any $m \geq 0$, we have*

$$\sum_{k\geq 0} h_k(-\mathbf{s})\frac{\partial^{m+1}\log Z^{\kappa}}{\partial t_0^m\partial t_k}(\mathbf{0};\mathbf{s}) = 0. \tag{B.12}$$

Comparing the coefficients of s_j in (B.12) we find

$$-\langle\tau_0^m\tau_j\rangle + \langle\tau_0^{m+1}\kappa_j\rangle = 0. \tag{B.13}$$

Substituting this expression into (B.11) and using (1.35),(6.3) we can solve out $A^{(j)}(z), B^{(j)}(z)$, e.g. in the case $j = 1$ we have

Lemma B.3. *$A^{(1)}(z)$ and $B^{(1)}(z)$ satisfy the following ODE system*

$$S_z^{WK}A^{(1)}(z) + B^{(1)}(z) = -\frac{1}{6}A^{(0)}(z) - \frac{z^3}{3}A^{(0)}(z) + \frac{z^2}{3}B^{(0)}(z), \tag{B.14}$$

$$S_z^{WK}B^{(1)}(z) + z^2A^{(1)}(z) = \frac{1}{6}B^{(0)}(z) - \frac{z^3}{3}B^{(0)}(z) + \frac{z^4}{3}A^{(0)}(z). \tag{B.15}$$

Moreover, the solution of this ODE system is unique under the boundary condition

$$A^{(1)}(z) = \mathcal{O}(z^{-1}), \qquad z \to \infty. \tag{B.16}$$

Recall that $A^{(0)}(z) = c(z)$, $B^{(0)}(z) = z\,q(z)$; we obtain the explicit solution of the ODE system (B.14),(B.15):

$$A^{(1)}(z) = \frac{1}{5}\sum_{g=1}^{\infty} g\,C_g\,z^{-3g+2}, \tag{B.17}$$

$$B^{(1)}(z) = \frac{1}{120}\sum_{g=0}^{\infty}(36g^2+48g-5)\,C_g\,z^{-3g}. \tag{B.18}$$

Similarly, we know from the system (B.8), (B.9) that for any partition λ,

$$S_z^{WK}A^{\lambda}(z) + B^{\lambda}(z) = \text{combinations of } A^{\mu}(z), B^{\mu}(z) \text{ with } |\mu| < |\lambda|, \tag{B.19}$$

$$S_z^{WK}B^{\lambda}(z) + z^2A^{\lambda}(z) = \text{combinations of } A^{\mu}(z), B^{\mu}(z) \text{ with } |\mu| < |\lambda|. \tag{B.20}$$

It is not difficult to see that the combination coefficients in the above formulæ can be obtained in a recursive procedure by applying Lemma B.2 and the *Main Theorem*. Details are omitted. So, in principle one can obtain all the initial data A^{λ}, B^{λ}, $|\lambda| > 0$ from (B.8), (B.9), (B.2) with the knowledge of $A^{(0)}, B^{(0)}$.

Now let $F_n^{\kappa}(z_1,\ldots,z_n;\mathbf{s}) := \nabla(z_1)\cdots\nabla(z_n)\log Z^{\kappa}(\mathbf{t};\mathbf{s})\,|_{\mathbf{t}=0}$ be the n-point generating function corresponding to the partition function Z^{κ}. We have for $n=1$,

$$\begin{aligned}
F_1^{\kappa}(z;\mathbf{s}) &= \frac{1}{4\,z}\left(-A(z;\mathbf{s})\,B_z(-z;\mathbf{s}) + B_z(z;\mathbf{s})\,A(-z;\mathbf{s}) + B(z;\mathbf{s})\,A_z(-z;\mathbf{s}) - A_z(z;\mathbf{s})\,B(-z;\mathbf{s})\right)\\
&= F_1(z) + \frac{s_1}{4\,z}\Big(-A^{WK}(z)\,B_z^{(1)}(-z) - A^{(1)}(z)\,B_z^{WK}(-z) + B_z^{WK}(z)\,A^{(1)}(-z) + B_z^{(1)}(z)\,A^{WK}(-z)\\
&\quad + B^{WK}(z)\,A_z^{(1)}(-z) + B^{(1)}(z)\,A_z^{WK}(-z) - A_z^{WK}(z)\,B^{(1)}(-z) - A_z^{(1)}(z)B^{WK}(-z)\Big)\\
&\quad + \text{higher order terms}\\
&= F_1(z) + s_1\sum_{g=1}^{\infty}\frac{(12g^2-12g+5)(6g-5)!!}{5\cdot 24^g\cdot g!}z^{-6g+4} + \text{ higher order terms.}
\end{aligned} \tag{B.21}$$

The last equality uses similar derivation as in Lemma 6.3. We read off from the above expression that

$$\langle\kappa_1\tau_{3g-3}\rangle_{g,1} = \frac{12g^2-12g+5}{5\cdot 24^g\cdot g!}, \qquad g \geq 1. \tag{B.22}$$

For $n \geq 2$, let $M^{\kappa}(z;\mathbf{s}) = \Theta(z;\mathbf{0};\mathbf{s})$ then we have

$$F_n^{\kappa}(z_1,\ldots,z_n;\mathbf{s}) = -\frac{1}{n}\sum_{r\in S_n}\frac{\mathrm{Tr}\left(M^{\kappa}(z_{r_1};\mathbf{s})\cdots M^{\kappa}(z_{r_n};\mathbf{s})\right)}{\prod_{j=1}^{n}(z_{r_j}^2 - z_{r_{j+1}}^2)} - \delta_{n,2}\frac{z_1^2+z_2^2}{(z_1^2-z_2^2)^2}, \tag{B.23}$$

Here the matrix-value function $M^{\kappa}(z;\mathbf{s})$ has the form

$$M^{\kappa}(z;\mathbf{s}) = M(z) + s_1\begin{pmatrix} M_{11}^{(1)}(z) & M_{12}^{(1)}(z) \\ M_{21}^{(1)}(z) & M_{22}^{(1)}(z)\end{pmatrix} + \text{higher order terms} \tag{B.24}$$

with

$$M_{11}^{(1)}(z) = -\frac{1}{2}\left(B^{WK}(z)\,A^{(1)}(-z) + B^{(1)}(z)\,A^{WK}(-z) + A^{WK}(z)\,B^{(1)}(-z) + A^{(1)}(z)\,B^{WK}(-z)\right), \quad \text{(B.25)}$$

$$M_{12}^{(1)}(z) = -A^{WK}(z)\,A^{(1)}(-z) - A^{(1)}(z)\,A^{WK}(-z), \quad \text{(B.26)}$$

$$M_{21}^{(1)}(z) = B^{WK}(z)\,B^{(1)}(-z) + B^{(1)}(z)\,B^{WK}(-z), \quad \text{(B.27)}$$

$$M_{22}^{(1)}(z) = -M_{12}^{(1)}(z). \quad \text{(B.28)}$$

C Tables of some intersection numbers

The nonzero two point intersection numbers $\langle \tau_k \tau_\ell \rangle$ with $2 \le k \le \ell \le 30$:

$$\langle \tau_3 \tau_2 \rangle = \frac{29}{5760}$$
$$\langle \tau_2 \tau_{12} \rangle = \frac{49}{955514880}$$
$$\langle \tau_2 \tau_{21} \rangle = \frac{677}{221911183337881600}$$
$$\langle \tau_2 \tau_{30} \rangle = \frac{53}{12148128371129859440640}$$
$$\langle \tau_3 \tau_{11} \rangle = \frac{307}{1672151040}$$
$$\langle \tau_3 \tau_{20} \rangle = \frac{4163}{221911183337881600}$$
$$\langle \tau_3 \tau_{29} \rangle = \frac{8087}{212592246494772540211200}$$
$$\langle \tau_4 \tau_{10} \rangle = \frac{29}{61931520}$$
$$\langle \tau_4 \tau_{19} \rangle = \frac{6319}{7397061112627200}$$
$$\langle \tau_4 \tau_{28} \rangle = \frac{17669}{70864082164924180070400}$$
$$\langle \tau_5 \tau_{12} \rangle = \frac{197089}{10594748989440}$$
$$\langle \tau_5 \tau_{21} \rangle = \frac{35759}{1318156290270167040}$$
$$\langle \tau_5 \tau_{30} \rangle = \frac{17981}{2499043550632428227788800}$$
$$\langle \tau_6 \tau_{14} \rangle = \frac{714211}{1101853894901760}$$
$$\langle \tau_6 \tau_{23} \rangle = \frac{697343}{913921694587315814400}$$
$$\langle {\tau_7}^2 \rangle = \frac{173}{116785152}$$
$$\langle \tau_7 \tau_{16} \rangle = \frac{1013239}{503704637669376000}$$
$$\langle \tau_7 \tau_{25} \rangle = \frac{439337}{227464955097287491584000}$$
$$\langle \tau_8 \tau_{12} \rangle = \frac{10964903}{5509269474508800}$$
$$\langle \tau_8 \tau_{21} \rangle = \frac{98324791}{1553666880798436888448000}$$
$$\langle \tau_8 \tau_{30} \rangle = \frac{227220239}{773204074565673293677854720000}$$
$$\langle \tau_9 \tau_{17} \rangle = \frac{2332617527}{2128822408786319769600}$$
$$\langle \tau_9 \tau_{26} \rangle = \frac{409854421}{3179843596698656402571264000}$$
$$\langle \tau_{10} \tau_{13} \rangle = \frac{143827807}{16347505058906112000}$$
$$\langle \tau_{10} \tau_{22} \rangle = \frac{8293101379}{247965234175430526763008000}$$
$$\langle \tau_{11} \tau_{12} \rangle = \frac{506527379}{4904251517671833600}$$
$$\langle \tau_{11} \tau_{21} \rangle = \frac{223913796019}{342192023162094126932951040}$$
$$\langle \tau_{11} \tau_{30} \rangle = \frac{265692198983}{523989695430585881802954296524800}$$

$$\langle \tau_2 \tau_6 \rangle = \frac{77}{414720}$$
$$\langle \tau_2 \tau_{15} \rangle = \frac{73}{137594142720}$$
$$\langle \tau_2 \tau_{24} \rangle = \frac{869}{47932956009824256000}$$
$$\langle \tau_3 \tau_5 \rangle = \frac{503}{1451520}$$
$$\langle \tau_3 \tau_{14} \rangle = \frac{2267}{963158999040}$$
$$\langle \tau_3 \tau_{23} \rangle = \frac{761}{5991619501228032000}$$
$$\langle {\tau_4}^2 \rangle = \frac{607}{1451520}$$
$$\langle \tau_4 \tau_{13} \rangle = \frac{7297}{963158999040}$$
$$\langle \tau_4 \tau_{22} \rangle = \frac{3973}{5991619501228032000}$$
$$\langle \tau_5 \tau_6 \rangle = \frac{487}{18579456}$$
$$\langle \tau_5 \tau_{15} \rangle = \frac{11519}{42378995957760}$$
$$\langle \tau_5 \tau_{24} \rangle = \frac{398477}{196844672680344944640 0}$$
$$\langle \tau_6 \tau_8 \rangle = \frac{2291}{1751777280}$$
$$\langle \tau_6 \tau_{17} \rangle = \frac{9119437}{1057779739105689600}$$
$$\langle \tau_6 \tau_{26} \rangle = \frac{671443}{1228310757525352454553 60}$$
$$\langle \tau_7 \tau_{10} \rangle = \frac{3777167}{68865868431360}$$
$$\langle \tau_7 \tau_{19} \rangle = \frac{42028829}{17136031773512171520 0}$$
$$\langle \tau_7 \tau_{28} \rangle = \frac{150998833}{11370648155377548436439040 00}$$
$$\langle \tau_8 \tau_{15} \rangle = \frac{9119281}{2335357865558016 00}$$
$$\langle \tau_8 \tau_{24} \rangle = \frac{604967959}{10440641438965495863705600}$$
$$\langle \tau_9 \tau_{11} \rangle = \frac{432827}{165278084235264}$$
$$\langle \tau_9 \tau_{20} \rangle = \frac{12683907763}{8855901220551090241536 00}$$
$$\langle \tau_9 \tau_{29} \rangle = \frac{20790655141}{2203631612512168886981885952000}$$
$$\langle \tau_{10} \tau_{16} \rangle = \frac{777536807}{4257644817572639539 20}$$
$$\langle \tau_{10} \tau_{25} \rangle = \frac{50685316067}{1558123362382341637259919360 0}$$
$$\langle \tau_{11} \tau_{15} \rangle = \frac{101418041}{38705861977933086720}$$
$$\langle \tau_{11} \tau_{24} \rangle = \frac{152055965491}{2108049254987873979822243840 0}$$
$$\langle \tau_{12} \tau_{14} \rangle = \frac{2515167869}{7741172395586617344 00}$$

$$\langle \tau_2 \tau_9 \rangle = \frac{149}{39813120}$$
$$\langle \tau_2 \tau_{18} \rangle = \frac{509}{1155790798848 00}$$
$$\langle \tau_2 \tau_{27} \rangle = \frac{31}{3286831269245091840 0}$$
$$\langle \tau_3 \tau_8 \rangle = \frac{947}{92897280}$$
$$\langle \tau_3 \tau_{17} \rangle = \frac{449}{19263179980800}$$
$$\langle \tau_3 \tau_{26} \rangle = \frac{1327}{1789497024366772224 00}$$
$$\langle \tau_4 \tau_7 \rangle = \frac{1781}{92897280}$$
$$\langle \tau_4 \tau_{16} \rangle = \frac{349}{3852635996160}$$
$$\langle \tau_4 \tau_{25} \rangle = \frac{7813}{1789497024366772224 00}$$
$$\langle \tau_5 \tau_9 \rangle = \frac{2351}{2627665920}$$
$$\langle \tau_5 \tau_{18} \rangle = \frac{739393}{2441030167166976 00}$$
$$\langle \tau_5 \tau_{27} \rangle = \frac{274679}{212592246494772540211200}$$
$$\langle \tau_6 \tau_{11} \rangle = \frac{492761}{13773173686272}$$
$$\langle \tau_6 \tau_{20} \rangle = \frac{3075329}{34272063547024343040}$$
$$\langle \tau_6 \tau_{29} \rangle = \frac{7677931}{2274129631075509687287808 00}$$
$$\langle \tau_7 \tau_{13} \rangle = \frac{6903563}{5509269474508800}$$
$$\langle \tau_7 \tau_{22} \rangle = \frac{2184989}{9139216945873158144000}$$
$$\langle \tau_8 \tau_9 \rangle = \frac{666643}{9837981204480}$$
$$\langle \tau_8 \tau_{18} \rangle = \frac{126087257}{2240865693459283968 00}$$
$$\langle \tau_8 \tau_{27} \rangle = \frac{8606940631}{1933010186414183234194636 8000}$$
$$\langle \tau_9 \tau_{14} \rangle = \frac{44636597}{7006073596674048 00}$$
$$\langle \tau_9 \tau_{23} \rangle = \frac{1588041583}{1062708146466130828984320 0}$$
$$\langle {\tau_{10}}^2 \rangle = \frac{474035}{165278084235264}$$
$$\langle \tau_{10} \tau_{19} \rangle = \frac{173346523879}{6199130854385763169075200}$$
$$\langle \tau_{10} \tau_{28} \rangle = \frac{408882825223}{1542542128758518220887 3201664000}$$
$$\langle \tau_{11} \tau_{18} \rangle = \frac{22610431297}{4768562195681356283904 00}$$
$$\langle \tau_{11} \tau_{27} \rangle = \frac{1226648541119}{1867287840076101004232019148 8000}$$
$$\langle \tau_{12} \tau_{17} \rangle = \frac{22308961201}{3179041463787570855936 00}$$

$$\langle \tau_{12}\tau_{20}\rangle = \frac{7702634988677}{6843840463241882538659020800}$$

$$\langle \tau_{12}\tau_{23}\rangle = \frac{532195893421}{37643736696212035353968640000}$$

$$\langle \tau_{12}\tau_{26}\rangle = \frac{545177136989}{377229866682040606915559424000 0}$$

$$\langle \tau_{12}\tau_{29}\rangle = \frac{32414448521083}{261994847715292940901477148262400 00}$$

$$\langle \tau_{13}{}^{2}\rangle = \frac{2701472413}{774117239558661734400}$$

$$\langle \tau_{13}\tau_{16}\rangle = \frac{4131287269}{4541487805410815508480 0}$$

$$\langle \tau_{13}\tau_{19}\rangle = \frac{3898864039667}{228128015441396084621967360 0}$$

$$\langle \tau_{13}\tau_{22}\rangle = \frac{926415021131}{37643736696212035353968640000}$$

$$\langle \tau_{13}\tau_{25}\rangle = \frac{1070162502821}{377229866682040606915559424000 0}$$

$$\langle \tau_{13}\tau_{28}\rangle = \frac{874463851081}{323450429278139433211700183040000}$$

$$\langle \tau_{14}\tau_{15}\rangle = \frac{427374701}{4128625277646195916800}$$

$$\langle \tau_{14}\tau_{18}\rangle = \frac{1209992413697}{526449266403221733743001600}$$

$$\langle \tau_{14}\tau_{21}\rangle = \frac{41072588543}{10755353341774867243991040 00}$$

$$\langle \tau_{14}\tau_{24}\rangle = \frac{110706467197}{22189992157767094524444672000 0}$$

$$\langle \tau_{14}\tau_{27}\rangle = \frac{212494716887933}{399886872828605015060149331558400 00}$$

$$\langle \tau_{14}\tau_{30}\rangle = \frac{6739031639609}{142089673254423807948645268979712000}$$

$$\langle \tau_{15}\tau_{17}\rangle = \frac{48139483517}{17548308880107391124766720}$$

$$\langle \tau_{15}\tau_{20}\rangle = \frac{398801587363}{752874733924240707079378 0000}$$

$$\langle \tau_{15}\tau_{23}\rangle = \frac{3124779323}{396249859960126687936512 0000}$$

$$\langle \tau_{15}\tau_{26}\rangle = \frac{761629810447}{807852268340616192040705720320 00}$$

$$\langle \tau_{15}\tau_{29}\rangle = \frac{41108093023159}{440477987088713804640800333837107200}$$

$$\langle \tau_{16}{}^{2}\rangle = \frac{450238303}{154746991888072232140800}$$

$$\langle \tau_{16}\tau_{19}\rangle = \frac{5450288228071}{828162207316664777787310080 00}$$

$$\langle \tau_{16}\tau_{22}\rangle = \frac{48954875677}{435874845956139356730163200 00}$$

$$\langle \tau_{16}\tau_{25}\rangle = \frac{4485153315589}{296212498391559270414925430784000}$$

$$\langle \tau_{16}\tau_{28}\rangle = \frac{4042295809491521}{242262892898792592552440183610408960 00}$$

$$\langle \tau_{17}\tau_{18}\rangle = \frac{16350865020703}{222966748123717440173506560000}$$

$$\langle \tau_{17}\tau_{21}\rangle = \frac{440593883623}{3051123921692975497111114240000}$$

$$\langle \tau_{17}\tau_{24}\rangle = \frac{5766625701953}{261363969169022885660228321280000}$$

$$\langle \tau_{17}\tau_{27}\rangle = \frac{1732412490886823}{637533928681033138295895220027392 0000}$$

$$\langle \tau_{17}\tau_{30}\rangle = \frac{41185538381700869}{148307023131086943614711208053676441600 00}$$

$$\langle \tau_{18}\tau_{20}\rangle = \frac{3072249243757}{183067435301578529826668544 0000}$$

$$\langle \tau_{18}\tau_{23}\rangle = \frac{5727662016017}{196022976876767161424517124096 0000}$$

$$\langle \tau_{18}\tau_{26}\rangle = \frac{6242927895721}{154553679680250457768701871521792 00}$$

$$\langle \tau_{18}\tau_{29}\rangle = \frac{20370144660007 0681}{444921069393260830844133624161029324800 00}$$

$$\langle \tau_{19}{}^{2}\rangle = \frac{3229800471133}{183067435301578529826668544 0000}$$

$$\langle \tau_{19}\tau_{22}\rangle = \frac{9970374606887}{283144299933108126131914014720000}$$

$$\langle \tau_{19}\tau_{25}\rangle = \frac{78779804364907}{143514131131661139356651737841664000}$$

$$\langle \tau_{19}\tau_{28}\rangle = \frac{4006128449215796443}{578397390211239080097373711409338122240000}$$

$$\langle \tau_{20}\tau_{21}\rangle = \frac{19697569368577}{509659739879594627037445226496000}$$

$$\langle \tau_{20}\tau_{24}\rangle = \frac{40350631517089}{590940539953898809115624802877440 00}$$

$$\langle \tau_{20}\tau_{27}\rangle = \frac{293131349971936519}{304419679058546884261775637583862169600 00}$$

$$\langle \tau_{20}\tau_{30}\rangle = \frac{271384734119406419}{242926903888720413640896958791922011340800 0}$$

$$\langle \tau_{21}\tau_{23}\rangle = \frac{40233333170437}{517072972459661457976171702517760 00}$$

$$\langle \tau_{21}\tau_{26}\rangle = \frac{5302117441409329}{430492475436328927238874639007481856000}$$

$$\langle \tau_{21}\tau_{29}\rangle = \frac{192493823039120893}{121463451944360206820448479395961005670 4000}$$

$$\langle \tau_{22}{}^{2}\rangle = \frac{210107406496993}{258536486229830728988085851258880000}$$

$$\langle \tau_{22}\tau_{25}\rangle = \frac{31223580485871083}{215246237718164463619437319503740928 0000}$$

$$\langle \tau_{22}\tau_{28}\rangle = \frac{4206346503347943 91}{202439086573933678034080798993268342784 0000}$$

$$\langle \tau_{23}\tau_{24}\rangle = \frac{1992994499234917}{126615433951861449187904305590435840000}$$

$$\langle \tau_{23}\tau_{27}\rangle = \frac{80547060703911827}{319640663011474228474864419463055278080000}$$

$$\langle \tau_{23}\tau_{30}\rangle = \frac{2893675174144460089}{878494034334232373673325110708163219292160000}$$

$$\langle \tau_{24}\tau_{26}\rangle = \frac{10228198184633927}{361613679366516298880654696766284759040 00}$$

$$\langle \tau_{24}\tau_{29}\rangle = \frac{100865248927375802843}{245978329613585064628531030998285701401804800 00}$$

$$\langle \tau_{25}{}^{2}\rangle = \frac{3543101331873197}{120537893122172099626884898922094919680 00}$$

$$\langle \tau_{25}\tau_{28}\rangle = \frac{1983683228896135764929}{418163160343094609868502752697085692383068160000}$$

$$\langle \tau_{26}\tau_{27}\rangle = \frac{1122839563530633017 29}{220085873864786636772896185630045101254246400 00}$$

$$\langle \tau_{26}\tau_{30}\rangle = \frac{2810189886598617689}{376527867321920687673803344419886684050554880 00}$$

$$\langle \tau_{27}\tau_{29}\rangle = \frac{771397123871360986807}{931906471621753701992663277439219543025123328000 0}$$

$$\langle \tau_{28}{}^{2}\rangle = \frac{7984636896204530513 59}{931906471621753701992663277439219543025123328000 0}$$

The nonzero three point intersection numbers $\langle \tau_{j_1}\tau_{j_2}\tau_{j_3}\rangle$ with $2 \le j_1 \le j_2 \le j_3 \le 22$:

$$\langle {\tau_2}^3\rangle = \frac{7}{240}$$
$$\langle {\tau_2}^2\tau_{14}\rangle = \frac{379}{9555148800}$$
$$\langle \tau_2\tau_3\tau_7\rangle = \frac{16243}{58060800}$$
$$\langle \tau_2\tau_3\tau_{19}\rangle = \frac{4889}{198135565516800}$$
$$\langle \tau_2\tau_4\tau_{12}\rangle = \frac{28279}{57330892800}$$
$$\langle \tau_2{\tau_5}^2\rangle = \frac{533}{967680}$$
$$\langle \tau_2\tau_5\tau_{17}\rangle = \frac{1979}{5503765708800}$$
$$\langle \tau_2\tau_6\tau_{13}\rangle = \frac{5818985}{110185389490176}$$
$$\langle \tau_2\tau_7\tau_9\rangle = \frac{11384627}{4099158835200}$$
$$\langle \tau_2\tau_7\tau_{21}\rangle = \frac{70348901}{1632003026048778240000}$$
$$\langle \tau_2\tau_8\tau_{17}\rangle = \frac{129856373}{1729060350760608000}$$
$$\langle \tau_2\tau_9\tau_{16}\rangle = \frac{2337414527}{1689541594274856960 0}$$
$$\langle \tau_2\tau_{10}\tau_{15}\rangle = \frac{2337411943}{10751628327203635200}$$
$$\langle \tau_2\tau_{11}\tau_{14}\rangle = \frac{105014281}{3583876109067878 40}$$
$$\langle \tau_2\tau_{12}\tau_{16}\rangle = \frac{558653463337}{5676859756763519385600 0}$$
$$\langle \tau_2\tau_{13}\tau_{18}\rangle = \frac{386519220661631}{1316123166008054334357504000}$$
$$\langle \tau_2\tau_{14}\tau_{20}\rangle = \frac{412571934041}{5228296763362782688051200 0}$$
$$\langle \tau_2\tau_{16}\tau_{18}\rangle = \frac{3273957891587}{2654366049091874287779840 00}$$
$$\langle \tau_2\tau_{18}\tau_{19}\rangle = \frac{46832106950971}{1307624537868418070190489600 00}$$
$$\langle \tau_2\tau_{21}\tau_{22}\rangle = \frac{47309184369850283}{2154470718581922741567382093824000 00}$$
$$\langle {\tau_3}^2\tau_{12}\rangle = \frac{3571489}{5618427494400}$$
$$\langle \tau_3\tau_4\tau_5\rangle = \frac{719}{829440}$$
$$\langle \tau_3\tau_4\tau_{17}\rangle = \frac{915071}{1618107118387200}$$
$$\langle \tau_3\tau_5\tau_{13}\rangle = \frac{14563891}{148326485852160}$$
$$\langle \tau_3{\tau_6}^2\rangle = \frac{1178689}{12262440960}$$
$$\langle \tau_3\tau_6\tau_{18}\rangle = \frac{9862412827}{1332802471273168896 00}$$
$$\langle \tau_3\tau_7\tau_{14}\rangle = \frac{8105473}{858587450572800}$$
$$\langle \tau_3\tau_8\tau_{13}\rangle = \frac{1645420379}{1021719066181632 00}$$
$$\langle \tau_3{\tau_9}^2\rangle = \frac{161547173}{317842469683200}$$

$$\langle {\tau_2}^2\tau_5\rangle = \frac{17}{5760}$$
$$\langle {\tau_2}^2\tau_{17}\rangle = \frac{7}{152882380 80}$$
$$\langle \tau_2\tau_3\tau_{10}\rangle = \frac{35521}{4180377600}$$
$$\langle \tau_2\tau_3\tau_{22}\rangle = \frac{35839}{166433875034112000}$$
$$\langle \tau_2\tau_4\tau_{15}\rangle = \frac{80897}{9631589990400}$$
$$\langle \tau_2\tau_5\tau_8\rangle = \frac{12491}{364953600}$$
$$\langle \tau_2\tau_5\tau_{20}\rangle = \frac{256801}{61025754179174400}$$
$$\langle \tau_2\tau_6\tau_{16}\rangle = \frac{831191}{858587450572800}$$
$$\langle \tau_2\tau_7\tau_{12}\rangle = \frac{261846317}{2754634737254400}$$
$$\langle \tau_2{\tau_8}^2\rangle = \frac{706933}{227731046400}$$
$$\langle \tau_2\tau_8\tau_{20}\rangle = \frac{1176392617}{1078935333887803392000}$$
$$\langle \tau_2\tau_9\tau_{19}\rangle = \frac{2604533679463}{1106987652568886280192000}$$
$$\langle \tau_2\tau_{10}\tau_{18}\rangle = \frac{2604532187203}{5960702744601695354880 00}$$
$$\langle \tau_2\tau_{11}\tau_{17}\rangle = \frac{38795383889}{5519169207964532736000}$$
$$\langle \tau_2\tau_{12}\tau_{19}\rangle = \frac{3478673146123919}{1710960115810470634664755 2000}$$
$$\langle \tau_2\tau_{13}\tau_{21}\rangle = \frac{1669477092113}{3136978058017669612830720 00}$$
$$\langle \tau_2\tau_{15}\tau_{16}\rangle = \frac{2651318414483}{6267253171466925401702400}$$
$$\langle \tau_2\tau_{16}\tau_{21}\rangle = \frac{163810545953}{6226783513659133667573760 00}$$
$$\langle \tau_2\tau_{18}\tau_{22}\rangle = \frac{171454834886651}{2227533828145081411876945920 0000}$$
$$\langle {\tau_3}^3\rangle = \frac{583}{96768}$$
$$\langle {\tau_3}^2\tau_{15}\rangle = \frac{1458581}{13484225986560 0}$$
$$\langle \tau_3\tau_4\tau_8\rangle = \frac{10723}{199065600}$$
$$\langle \tau_3\tau_4\tau_{20}\rangle = \frac{1541879}{233007425047756800}$$
$$\langle \tau_3\tau_5\tau_{16}\rangle = \frac{4575643}{2542739757465600}$$
$$\langle \tau_3\tau_6\tau_9\rangle = \frac{1243987}{208684449792}$$
$$\langle \tau_3\tau_6\tau_{21}\rangle = \frac{591294929}{6397451862111210700 80}$$
$$\langle \tau_3\tau_7\tau_{17}\rangle = \frac{10136254319}{5553343630304870400 0}$$
$$\langle \tau_3\tau_8\tau_{16}\rangle = \frac{596252071}{1586669608658534400}$$
$$\langle \tau_3\tau_9\tau_{12}\rangle = \frac{14029444187}{6130314397089792 00}$$

$$\langle {\tau_2}^2\tau_8\rangle = \frac{53}{460800}$$
$$\langle {\tau_2}^2\tau_{20}\rangle = \frac{139}{33022594252800}$$
$$\langle \tau_2\tau_3\tau_{13}\rangle = \frac{9433}{57330892800}$$
$$\langle \tau_2\tau_4\tau_6\rangle = \frac{1081}{2322432}$$
$$\langle \tau_2\tau_4\tau_{18}\rangle = \frac{21179}{198135565516800}$$
$$\langle \tau_2\tau_5\tau_{11}\rangle = \frac{5893}{5255331840}$$
$$\langle \tau_2\tau_6\tau_7\rangle = \frac{195983}{4379443200}$$
$$\langle \tau_2\tau_6\tau_{19}\rangle = \frac{126347401}{95200176519512064 00}$$
$$\langle \tau_2\tau_7\tau_{15}\rangle = \frac{1828589}{858587450572800}$$
$$\langle \tau_2\tau_8\tau_{11}\rangle = \frac{39617}{283398635520}$$
$$\langle \tau_2\tau_9\tau_{10}\rangle = \frac{13984247}{82639042117632}$$
$$\langle \tau_2\tau_9\tau_{22}\rangle = \frac{117449747941}{3795386237379038674944000}$$
$$\langle \tau_2\tau_{10}\tau_{21}\rangle = \frac{2466443967737}{3719478512631457901445120 0}$$
$$\langle \tau_2\tau_{11}\tau_{20}\rangle = \frac{17676185055899}{1425800096508725528888729600}$$
$$\langle \tau_2\tau_{12}\tau_{22}\rangle = \frac{5008431389959}{1568489029008834806415360000}$$
$$\langle {\tau_2\tau_{14}}^2\rangle = \frac{11058913867}{8601302661762908160 00}$$
$$\langle \tau_2\tau_{15}\tau_{19}\rangle = \frac{3273957903823}{3136978058017669612830720 00}$$
$$\langle {\tau_2\tau_{17}}^2\rangle = \frac{3364901200937}{2580633658883932222423040000}$$
$$\langle \tau_2\tau_{19}\tau_{21}\rangle = \frac{514364504566573}{5791587953177211670880059392000 0}$$
$$\langle {\tau_3}^2\tau_6\rangle = \frac{1997}{3317760}$$
$$\langle {\tau_3}^2\tau_{18}\rangle = \frac{2671901}{19417285420646400}$$
$$\langle \tau_3\tau_4\tau_{11}\rangle = \frac{247823}{140460687360}$$
$$\langle \tau_3\tau_5\tau_7\rangle = \frac{1276397}{15328051200}$$
$$\langle \tau_3\tau_5\tau_{19}\rangle = \frac{63219559}{2563081675525324800}$$
$$\langle \tau_3\tau_6\tau_{12}\rangle = \frac{120998503}{593305943408640}$$
$$\langle \tau_3\tau_7\tau_8\rangle = \frac{6564751}{869518540800}$$
$$\langle \tau_3\tau_7\tau_{20}\rangle = \frac{3026847413}{1142402118234144768000}$$
$$\langle \tau_3\tau_8\tau_{19}\rangle = \frac{124100949853}{1942083600998046105600 0}$$
$$\langle \tau_3\tau_9\tau_{15}\rangle = \frac{10732557077}{1644366685337026560 0}$$

$$\langle {\tau_2}^2\tau_{11}\rangle = \frac{257}{99532800}$$
$$\langle \tau_2\tau_3\tau_4\rangle = \frac{1121}{241920}$$
$$\langle \tau_2\tau_3\tau_{16}\rangle = \frac{22073}{9631589990400}$$
$$\langle \tau_2\tau_4\tau_9\rangle = \frac{11827}{597196800}$$
$$\langle \tau_2\tau_4\tau_{21}\rangle = \frac{35831}{3328677500682240 0}$$
$$\langle \tau_2\tau_5\tau_{14}\rangle = \frac{16721}{706316599296}$$
$$\langle \tau_2\tau_6\tau_{10}\rangle = \frac{1626457}{819831767040}$$
$$\langle \tau_2\tau_6\tau_{22}\rangle = \frac{164148389}{1142402118234144768000}$$
$$\langle \tau_2\tau_7\tau_{18}\rangle = \frac{126346169}{3661545250750464000}$$
$$\langle \tau_2\tau_8\tau_{14}\rangle = \frac{9447803}{2432664443289600}$$
$$\langle \tau_2\tau_9\tau_{13}\rangle = \frac{259565549}{43787959979212800}$$
$$\langle \tau_2\tau_{10}\tau_{12}\rangle = \frac{333726503}{43787959979212800}$$
$$\langle \tau_2{\tau_{11}}^2\rangle = \frac{604579}{72979933298688}$$
$$\langle \tau_2\tau_{12}\tau_{13}\rangle = \frac{73089896617}{2150325665440727040 00}$$
$$\langle \tau_2\tau_{13}\tau_{15}\rangle = \frac{62072602013}{5160781597057744896000}$$
$$\langle \tau_2\tau_{14}\tau_{17}\rangle = \frac{13698478951523}{3655897683355706484326400 0}$$
$$\langle \tau_2\tau_{15}\tau_{22}\rangle = \frac{382224608021}{1981249299800633439682560000}$$
$$\langle \tau_2\tau_{17}\tau_{20}\rangle = \frac{130441731463}{4035878203297586636390400 00}$$
$$\langle {\tau_2\tau_{20}}^2\rangle = \frac{89909242721189}{9652646588628686118133432320000}$$
$$\langle {\tau_3}^2\tau_9\rangle = \frac{213631}{8360755200}$$
$$\langle {\tau_3}^2\tau_{21}\rangle = \frac{645587}{4660148500955136 00}$$
$$\langle \tau_3\tau_4\tau_{14}\rangle = \frac{502151}{13484225986560}$$
$$\langle \tau_3\tau_5\tau_{10}\rangle = \frac{518267}{140460687360}$$
$$\langle \tau_3\tau_5\tau_{22}\rangle = \frac{82123427}{3075698010630389760 00}$$
$$\langle \tau_3\tau_6\tau_{15}\rangle = \frac{10981843}{2404044861603840}$$
$$\langle \tau_3\tau_7\tau_{11}\rangle = \frac{126034451}{370816214630400}$$
$$\langle \tau_3\tau_8\tau_{10}\rangle = \frac{170518829}{370816214630400}$$
$$\langle \tau_3\tau_8\tau_{22}\rangle = \frac{274189207879}{3262700449676717457408000}$$
$$\langle \tau_3\tau_9\tau_{18}\rangle = \frac{117569418301}{8963462773837135872000}$$

$$\langle \tau_3\tau_9\tau_{21}\rangle = \frac{2961244699003}{14877914050525831605780480}$$
$$\langle \tau_3\tau_{10}\tau_{20}\rangle = \frac{15158746939103}{3719478512631457901445120 0}$$
$$\langle \tau_3\tau_{11}\tau_{22}\rangle = \frac{272315357551}{23868311311004007923712000}$$
$$\langle \tau_3\tau_{12}\tau_{21}\rangle = \frac{19606706386613}{954732452440160316948480000}$$
$$\langle \tau_3\tau_{14}\tau_{16}\rangle = \frac{11747546324191}{626725317146692540170240 0}$$
$$\langle \tau_3\tau_{15}\tau_{18}\rangle = \frac{610318748197}{1049156541143033315328000 0}$$
$$\langle \tau_3\tau_{17}\tau_{19}\rangle = \frac{76848896914045 61}{40579948158516574111578193920 00}$$
$$\langle \tau_3\tau_{19}\tau_{20}\rangle = \frac{243373186276705439}{44595227239464529865776457318400 00}$$
$$\langle {\tau_4}^2\tau_7\rangle = \frac{15727}{154828800}$$
$$\langle {\tau_4}^2\tau_{19}\rangle = \frac{7022437}{233007425047756800}$$
$$\langle \tau_4\tau_5\tau_{12}\rangle = \frac{43670197}{148326485852160}$$
$$\langle \tau_4\tau_6\tau_8\rangle = \frac{437341}{34780741632}$$
$$\langle \tau_4\tau_6\tau_{20}\rangle = \frac{3103951}{7030166881440890 88}$$
$$\langle \tau_4\tau_7\tau_{16}\rangle = \frac{160854971}{226667086951219200}$$
$$\langle \tau_4\tau_8\tau_{12}\rangle = \frac{54280766647}{11238909072799795200}$$
$$\langle \tau_4\tau_9\tau_{11}\rangle = \frac{10713327791}{168583645919969280}$$
$$\langle {\tau_4\tau_{10}}^2\rangle = \frac{27378271519}{393361840479928320}$$
$$\langle \tau_4\tau_{10}\tau_{22}\rangle = \frac{11829432500983}{40576129228706813470310 4000}$$
$$\langle \tau_4\tau_{11}\tau_{21}\rangle = \frac{925781758043}{16230451691482725388124160}$$
$$\langle {\tau_4\tau_{13}}^2\rangle = \frac{326478195847}{190134058838969 5488000}$$
$$\langle \tau_4\tau_{14}\tau_{15}\rangle = \frac{3915350347109}{56975028831517503650 18400}$$
$$\langle \tau_4\tau_{15}\tau_{20}\rangle = \frac{62473991504027}{10540246274939369899111219200}$$
$$\langle \tau_4\tau_{17}\tau_{18}\rangle = \frac{2561433648422567}{31215344737320441624290918 4000}$$
$$\langle \tau_4\tau_{19}\tau_{22}\rangle = \frac{502288956613107577}{81721303118624655714624838041600000}$$
$$\langle {\tau_5}^2\tau_8\rangle = \frac{263}{17694720}$$
$$\langle {\tau_5}^2\tau_{20}\rangle = \frac{16813381}{32221598206604083 20}$$
$$\langle \tau_5\tau_6\tau_{16}\rangle = \frac{43871981}{45333417390243840}$$
$$\langle \tau_5\tau_7\tau_{12}\rangle = \frac{4934925797}{661112336941056 00}$$
$$\langle \tau_5{\tau_8}^2\rangle = \frac{5452297}{2942985830400}$$
$$\langle \tau_5\tau_8\tau_{20}\rangle = \frac{280687170067}{17090335688782805729280 0}$$

$$\langle \tau_3\tau_{10}\tau_{11}\rangle = \frac{2922785743}{107280501949071360}$$
$$\langle \tau_3\tau_{11}\tau_{13}\rangle = \frac{13983412199}{11510566797359185920}$$
$$\langle {\tau_3\tau_{12}}^2\rangle = \frac{3020419917227}{230211335947187184000}$$
$$\langle \tau_3\tau_{13}\tau_{14}\rangle = \frac{14476869931}{271620084055670784000}$$
$$\langle \tau_3\tau_{14}\tau_{19}\rangle = \frac{3153313491137}{68195175174297165496320000}$$
$$\langle \tau_3\tau_{15}\tau_{21}\rangle = \frac{1046156286335}{84321970199514959192889 7536}$$
$$\langle \tau_3\tau_{17}\tau_{22}\rangle = \frac{11634118724423347}{28586684127861878119084 7426400000}$$
$$\langle \tau_3\tau_{20}\tau_{22}\rangle = \frac{110264023018328843}{81721303118624655714624838041600000}$$
$$\langle {\tau_4}^2\tau_{10}\rangle = \frac{632767}{140460687360}$$
$$\langle {\tau_4}^2\tau_{22}\rangle = \frac{9123013}{27960891005730816000}$$
$$\langle \tau_4\tau_5\tau_{15}\rangle = \frac{3354359}{508547951493120}$$
$$\langle \tau_4\tau_6\tau_{11}\rangle = \frac{41992117}{74163242926080}$$
$$\langle {\tau_4\tau_7}^2\rangle = \frac{1769879}{124216934400}$$
$$\langle \tau_4\tau_7\tau_{19}\rangle = \frac{1060509119}{878770860180111360 00}$$
$$\langle \tau_4\tau_8\tau_{15}\rangle = \frac{198704189}{144242691696230400}$$
$$\langle \tau_4\tau_9\tau_{14}\rangle = \frac{2639929909}{11745476323835904 00}$$
$$\langle \tau_4\tau_{10}\tau_{13}\rangle = \frac{178634783759}{57552833986795929600}$$
$$\langle \tau_4\tau_{11}\tau_{12}\rangle = \frac{209702107861}{57552833986795929600}$$
$$\langle \tau_4\tau_{12}\tau_{14}\rangle = \frac{303962593411}{19013405883896954880 00}$$
$$\langle \tau_4\tau_{13}\tau_{16}\rangle = \frac{12616127590387}{20890843904889751339008 00}$$
$$\langle \tau_4\tau_{14}\tau_{18}\rangle = \frac{125069214236137}{62424814198010482262016 0000}$$
$$\langle {\tau_4\tau_{16}}^2\rangle = \frac{8293119762493}{32698712198957871661056000}$$
$$\langle \tau_4\tau_{17}\tau_{21}\rangle = \frac{11633407002466331}{57173368255723756238174945280000}$$
$$\langle \tau_4\tau_{20}\tau_{21}\rangle = \frac{15751221641063269}{23348943748178473061321382297600 00}$$
$$\langle {\tau_5}^2\tau_{11}\rangle = \frac{4135963}{6180270243840}$$
$$\langle \tau_5\tau_6\tau_7\rangle = \frac{3380051}{173903708160}$$
$$\langle \tau_5\tau_6\tau_{19}\rangle = \frac{8272260727}{50265693203023697920}$$
$$\langle \tau_5\tau_7\tau_{15}\rangle = \frac{3378105877}{15866696068658534400}$$
$$\langle \tau_5\tau_8\tau_{11}\rangle = \frac{1028118803}{9365758106664960}$$
$$\langle \tau_5\tau_9\tau_{10}\rangle = \frac{22402132091}{16858364591996928 0}$$

$$\langle \tau_3\tau_{10}\tau_{14}\rangle = \frac{11090288141}{11510566797359185920}$$
$$\langle \tau_3\tau_{11}\tau_{16}\rangle = \frac{210148115}{597564184922475724 8}$$
$$\langle \tau_3\tau_{12}\tau_{15}\rangle = \frac{25217780039}{54324016811134156 8000}$$
$$\langle \tau_3\tau_{13}\tau_{17}\rangle = \frac{340678837517771}{21935386100134238905958 4000}$$
$$\langle \tau_3\tau_{14}\tau_{22}\rangle = \frac{45042839922089}{52701231374696848494955560 96000}$$
$$\langle \tau_3\tau_{16}\tau_{17}\rangle = \frac{627272037989}{96172682938111138723840000}$$
$$\langle {\tau_3\tau_{18}}^2\rangle = \frac{7477189991322341}{37458413684784529949149102080 00}$$
$$\langle {\tau_3\tau_{21}}^2\rangle = \frac{46157032898850703}{32688521247449862285849935216640000}$$
$$\langle {\tau_4}^2\tau_{13}\rangle = \frac{323437}{2696845197312}$$
$$\langle \tau_4\tau_5\tau_6\rangle = \frac{424987}{3065610240}$$
$$\langle \tau_4\tau_5\tau_{18}\rangle = \frac{273887629}{25630816755253248 00}$$
$$\langle \tau_4\tau_6\tau_{14}\rangle = \frac{207975043}{13222246738821120}$$
$$\langle \tau_4\tau_7\tau_{10}\rangle = \frac{321920519}{3708162140304 00}$$
$$\langle \tau_4\tau_7\tau_{22}\rangle = \frac{10153851527}{63974518621112107008000}$$
$$\langle \tau_4\tau_8\tau_{18}\rangle = \frac{41359951891}{14939104623061893120 00}$$
$$\langle \tau_4\tau_9\tau_{17}\rangle = \frac{40271556439}{74695523115309465 6000}$$
$$\langle \tau_4\tau_{10}\tau_{16}\rangle = \frac{18793373881}{20914746472286650368 0}$$
$$\langle \tau_4\tau_{11}\tau_{15}\rangle = \frac{490262215}{3802681176779390976}$$
$$\langle \tau_4\tau_{12}\tau_{17}\rangle = \frac{1021906469984573}{21935386100134238905958 4000}$$
$$\langle \tau_4\tau_{13}\tau_{19}\rangle = \frac{1209002367692141}{81152258457413626940620 80000}$$
$$\langle \tau_4\tau_{14}\tau_{21}\rangle = \frac{45039389150641}{10540246274939369899111219200}$$
$$\langle \tau_4\tau_{16}\tau_{19}\rangle = \frac{4269056056230239}{57971354512166534445117056000}$$
$$\langle \tau_4\tau_{18}\tau_{20}\rangle = \frac{81119432622320849}{34304020953434253742904967 1680000}$$
$$\langle {\tau_4\tau_{22}}^2\rangle = \frac{24449165035836451}{14113654335127235825665232934912000 00}$$
$$\langle {\tau_5}^2\tau_{14}\rangle = \frac{1575647}{84757991915520}$$
$$\langle \tau_5\tau_6\tau_{10}\rangle = \frac{87811657}{74163242926080}$$
$$\langle \tau_5\tau_6\tau_{22}\rangle = \frac{3655470977}{16889272915973596250112}$$
$$\langle \tau_5\tau_7\tau_{18}\rangle = \frac{5908719973}{13809256374258892800 0}$$
$$\langle \tau_5\tau_8\tau_{14}\rangle = \frac{48889693}{12592615941734400}$$
$$\langle \tau_5\tau_9\tau_{13}\rangle = \frac{76561282313}{12920023956219494400}$$

$$\langle \tau_3\tau_{10}\tau_{17}\rangle = \frac{120835139197}{52286866180716625920 00}$$
$$\langle \tau_3\tau_{11}\tau_{19}\rangle = \frac{621508694340991}{85548005790523531733237760 0}$$
$$\langle \tau_3\tau_{12}\tau_{18}\rangle = \frac{29832419784677 27}{26322463320161086687150080 00}$$
$$\langle \tau_3\tau_{13}\tau_{20}\rangle = \frac{2230392447653}{68195175174297165496320000}$$
$$\langle {\tau_3\tau_{15}}^2\rangle = \frac{2273718760099}{11395005766303500730368 00}$$
$$\langle \tau_3\tau_{16}\tau_{20}\rangle = \frac{937181668915301}{57971354512166534445117056000}$$
$$\langle \tau_3\tau_{18}\tau_{21}\rangle = \frac{33959049274063673}{68608041906868507485809934336 0000}$$
$$\langle {\tau_4}^3\rangle = \frac{175}{165888}$$
$$\langle {\tau_4}^2\tau_{16}\rangle = \frac{169399}{77052719923200}$$
$$\langle \tau_4\tau_5\tau_9\rangle = \frac{34525}{4013162496}$$
$$\langle \tau_4\tau_5\tau_{21}\rangle = \frac{16421903}{12302792042521559040}$$
$$\langle \tau_4\tau_6\tau_{17}\rangle = \frac{3377997577}{11106687260609740800}$$
$$\langle \tau_4\tau_7\tau_{13}\rangle = \frac{670124677}{22037077898035200}$$
$$\langle \tau_4\tau_8\tau_9\rangle = \frac{8115953}{7567677849600}$$
$$\langle \tau_4\tau_8\tau_{21}\rangle = \frac{54830864083}{13050801798706869829632 0}$$
$$\langle \tau_4\tau_9\tau_{20}\rangle = \frac{1010454846697}{10627081464661308289843 20}$$
$$\langle \tau_4\tau_{10}\tau_{19}\rangle = \frac{23015905616471}{12398261708771526338150400}$$
$$\langle \tau_4\tau_{11}\tau_{18}\rangle = \frac{207143195266297}{65806158300402716717875200}$$
$$\langle \tau_4\tau_{12}\tau_{20}\rangle = \frac{796172328305677}{81152258457413626940620 80000}$$
$$\langle \tau_4\tau_{13}\tau_{22}\rangle = \frac{145126919082839}{52701231374696849495556096000}$$
$$\langle \tau_4\tau_{15}\tau_{17}\rangle = \frac{24879359891489}{10404135699668413710336 0000}$$
$$\langle \tau_4\tau_{16}\tau_{22}\rangle = \frac{15958034270869}{10083486464854277996150784 0000}$$
$$\langle {\tau_4\tau_{19}}^2\rangle = \frac{123181360481069857}{49550252488293922073084952576 0000}$$
$$\langle {\tau_5}^3\rangle = \frac{41873}{25546752 0}$$
$$\langle {\tau_5}^2\tau_{17}\rangle = \frac{8530549}{23732237736345600}$$
$$\langle \tau_5\tau_6\tau_{13}\rangle = \frac{548335477}{13222246738821120}$$
$$\langle \tau_5\tau_7\tau_9\rangle = \frac{12544139}{7567677849600}$$
$$\langle \tau_5\tau_7\tau_{21}\rangle = \frac{274159253063}{42223182289939906252800}$$
$$\langle \tau_5\tau_8\tau_{17}\rangle = \frac{6072863851}{65210377322889216000}$$
$$\langle \tau_5\tau_9\tau_{16}\rangle = \frac{56381768141}{32866030170736164864 0}$$

$$\langle \tau_5\tau_9\tau_{19}\rangle = \frac{207147188280589}{58448948055637195594137600}$$

$$\langle \tau_5\tau_{10}\tau_{15}\rangle = \frac{1025122387}{38026811767790976}$$

$$\langle \tau_5{\tau_{11}}^2\rangle = \frac{8737971317}{1055135289757925376}$$

$$\langle \tau_5\tau_{11}\tau_{20}\rangle = \frac{364917172905469}{16365705455578414766358528000}$$

$$\langle \tau_5\tau_{12}\tau_{19}\rangle = \frac{359078501200526983}{9819423273347048859815116800000}$$

$$\langle \tau_5\tau_{13}\tau_{18}\rangle = \frac{39897610190525287}{75534025179592683537039360000}$$

$$\langle \tau_5\tau_{14}\tau_{17}\rangle = \frac{52370214509563}{777099024481406209228800000}$$

$$\langle \tau_5\tau_{15}\tau_{19}\rangle = \frac{232859834319137}{10540246274939369899111219200}$$

$$\langle \tau_5\tau_{16}\tau_{21}\rangle = \frac{151235358464002367}{233594047444814204058829062144000}$$

$$\langle \tau_5\tau_{18}\tau_{19}\rangle = \frac{43236962777465398453}{490547499634109828523541030502400000}$$

$$\langle \tau_5{\tau_{20}}^2\rangle = \frac{790190487014602409}{29964477810162373762029107281920000}$$

$$\langle {\tau_6}^2\tau_9\rangle = \frac{1052741}{550376570880}$$

$$\langle {\tau_6}^2\tau_{18}\rangle = \frac{99268122211}{2010627728092094791680}$$

$$\langle \tau_6\tau_7\tau_{11}\rangle = \frac{73003477}{508547951493120}$$

$$\langle \tau_6\tau_7\tau_{20}\rangle = \frac{259097321723}{120637663685525687500800}$$

$$\langle \tau_6\tau_8\tau_{16}\rangle = \frac{8402066105}{3351046213486824652 8}$$

$$\langle {\tau_6\tau_9}^2\rangle = \frac{144611581}{673660922757120}$$

$$\langle \tau_6\tau_9\tau_{18}\rangle = \frac{130830697680127}{12305041695923620125081600}$$

$$\langle \tau_6\tau_{10}\tau_{14}\rangle = \frac{205629494617}{319872593105560535040}$$

$$\langle \tau_6\tau_{11}\tau_{13}\rangle = \frac{259272029251}{319872593105560535040}$$

$$\langle \tau_6\tau_{11}\tau_{22}\rangle = \frac{85184312423297}{65532835535492604155343667 20}$$

$$\langle \tau_6\tau_{12}\tau_{18}\rangle = \frac{43089630285107502 43}{392776930933881954392604672 0000}$$

$$\langle \tau_6\tau_{13}\tau_{17}\rangle = \frac{37851763107632743}{25178008393197561179013120000}$$

$$\langle \tau_6\tau_{14}\tau_{19}\rangle = \frac{69048249744375 13}{13106567107098520831068734400 0}$$

$$\langle \tau_6\tau_{15}\tau_{18}\rangle = \frac{1336417742448461}{20163949395536185893951897600}$$

$$\langle \tau_6\tau_{16}\tau_{20}\rangle = \frac{98847695400150793 79}{46158183775095286722024622679654 40}$$

$$\langle {\tau_6\tau_{18}}^2\rangle = \frac{19716107920024747086 71}{74563219944384693935578236636364 8000}$$

$$\langle \tau_6\tau_{20}\tau_{22}\rangle = \frac{84830653301124489 47}{36189356426209008827431283117260 80000}$$

$$\langle {\tau_7}^2\tau_{10}\rangle = \frac{186559673}{847579919155200}$$

$$\langle {\tau_7}^2\tau_{19}\rangle = \frac{393442486447}{67020924269736493056000}$$

$$\langle \tau_7\tau_8\tau_{12}\rangle = \frac{35336068039}{2266670869512192000}$$

$$\langle \tau_7\tau_8\tau_{21}\rangle = \frac{780754789932437}{31993108409401412325212160 00}$$

$$\langle \tau_7\tau_9\tau_{17}\rangle = \frac{134464346825603}{51271007066348417187840000}$$

$$\langle \tau_5\tau_9\tau_{22}\rangle = \frac{390376552056761}{70138737666764634712965120 00}$$

$$\langle \tau_5\tau_{10}\tau_{18}\rangle = \frac{41429423338949}{62945020982993902947532 80}$$

$$\langle \tau_5\tau_{11}\tau_{14}\rangle = \frac{12665466509}{34857910787144417280}$$

$$\langle \tau_5\tau_{12}\tau_{13}\rangle = \frac{8815163665079}{20914746472286650368000}$$

$$\langle \tau_5\tau_{12}\tau_{22}\rangle = \frac{3918464405395483}{57971354512166534445111 7056000}$$

$$\langle \tau_5\tau_{13}\tau_{21}\rangle = \frac{1306154784535829}{11594270902433306889022341 1200}$$

$$\langle \tau_5\tau_{14}\tau_{20}\rangle = \frac{322785380359679}{19323784837388844815037235200}$$

$$\langle \tau_5\tau_{15}\tau_{22}\rangle = \frac{4582889655795079}{96526465886286861181334432320000}$$

$$\langle {\tau_5\tau_{17}}^2\rangle = \frac{526521959129899}{19076044006140269881511116800}$$

$$\langle \tau_5\tau_{18}\tau_{22}\rangle = \frac{15068748821188750 21}{69148794946528554835451786035200000}$$

$$\langle \tau_5\tau_{21}\tau_{22}\rangle = \frac{37150181583737429}{52422144673329733066708651868160000}$$

$$\langle {\tau_6}^2\tau_{12}\rangle = \frac{31856771}{369853055631360}$$

$$\langle {\tau_6}^2\tau_{21}\rangle = \frac{253072809667}{33778545831947192500 2240}$$

$$\langle \tau_6\tau_7\tau_{14}\rangle = \frac{16111200341}{3173339217317068800}$$

$$\langle \tau_6\tau_8\tau_{10}\rangle = \frac{1679089523}{8645315175383040}$$

$$\langle \tau_6\tau_8\tau_{19}\rangle = \frac{10623001356787}{20508402826539366875136 00}$$

$$\langle \tau_6\tau_9\tau_{12}\rangle = \frac{636057301397}{5168009582487797760 0}$$

$$\langle \tau_6\tau_9\tau_{21}\rangle = \frac{14053624463775079}{72944287173435220101483724800}$$

$$\langle \tau_6\tau_{10}\tau_{17}\rangle = \frac{26892966289403}{14355881978577556812595 20}$$

$$\langle \tau_6\tau_{11}\tau_{16}\rangle = \frac{134464848878407}{47169326501040543812812 80}$$

$$\langle {\tau_6\tau_{12}}^2\rangle = \frac{56002776712291}{63974518621112107008000}$$

$$\langle \tau_6\tau_{12}\tau_{21}\rangle = \frac{6133270572795403}{26213134214197041662137466 8800}$$

$$\langle \tau_6\tau_{13}\tau_{20}\rangle = \frac{488390056473327 7}{13106567107098520831068733440 0}$$

$$\langle \tau_6\tau_{14}\tau_{22}\rangle = \frac{11877048677412430103}{10490496312521656073187414245376000}$$

$$\langle \tau_6\tau_{15}\tau_{21}\rangle = \frac{275854034160526373}{16784794100034649717099862792 6016}$$

$$\langle \tau_6\tau_{17}\tau_{19}\rangle = \frac{20263777566231577640 39}{80776821606416751763543089689 3952000}$$

$$\langle \tau_6\tau_{18}\tau_{21}\rangle = \frac{54247607002735273571}{71914746744389697028869857476608 0000}$$

$$\langle {\tau_6\tau_{21}}^2\rangle = \frac{3551050603479151147}{14475742570483603530972513246904320 00}$$

$$\langle {\tau_7}^2\tau_{13}\rangle = \frac{155739672533}{158666960865853440 00}$$

$$\langle {\tau_7}^2\tau_{22}\rangle = \frac{2478585038729}{26884965049917153214464000}$$

$$\langle \tau_7\tau_8\tau_{15}\rangle = \frac{210050295797}{38080070607804825 6000}$$

$$\langle \tau_7\tau_9\tau_{11}\rangle = \frac{26502074641}{1292002395621949440}$$

$$\langle \tau_7\tau_9\tau_{20}\rangle = \frac{1007173881978686 81}{18236071793358805025370931200 0}$$

$$\langle \tau_5\tau_{10}\tau_{12}\rangle = \frac{4823355455459}{6330811738547552256 00}$$

$$\langle \tau_5\tau_{10}\tau_{21}\rangle = \frac{1171129439636251}{98194232733470488598151168 00}$$

$$\langle \tau_5\tau_{11}\tau_{17}\rangle = \frac{212901240024101}{20107437258456385663795200}$$

$$\langle \tau_5\tau_{12}\tau_{16}\rangle = \frac{1021925948176829}{68939784886136179418726400}$$

$$\langle \tau_5\tau_{13}\tau_{15}\rangle = \frac{113547322348489}{62672531714669254017024 00}$$

$$\langle {\tau_5\tau_{14}}^2\rangle = \frac{20229696415813}{10445421952444875669504 00}$$

$$\langle \tau_5\tau_{15}\tau_{16}\rangle = \frac{273676602813451}{35968583418853658827161 6000}$$

$$\langle \tau_5\tau_{16}\tau_{18}\rangle = \frac{12807290860841347}{49052684587217836838171443 2000}$$

$$\langle \tau_5\tau_{17}\tau_{20}\rangle = \frac{10838534034576022 51}{13626319434280828570009836195840000}$$

$$\langle \tau_5\tau_{19}\tau_{21}\rangle = \frac{45206246459050251 23}{17978686686097424257217464369152 0000}$$

$$\langle {\tau_6}^3\rangle = \frac{3120571}{139122966528}$$

$$\langle {\tau_6}^2\tau_{15}\rangle = \frac{283484827}{115394153356984320}$$

$$\langle \tau_6\tau_7\tau_8\rangle = \frac{7778203}{3210529996800}$$

$$\langle \tau_6\tau_7\tau_{17}\rangle = \frac{510124481959}{41888077668585308160 00}$$

$$\langle \tau_6\tau_8\tau_{13}\rangle = \frac{3926272391}{4533341739024384 00}$$

$$\langle \tau_6\tau_8\tau_{22}\rangle = \frac{260252249259347}{31993108409401412325212160 00}$$

$$\langle \tau_6\tau_9\tau_{15}\rangle = \frac{39799290257}{91392169458731581440}$$

$$\langle \tau_6\tau_{10}\tau_{11}\rangle = \frac{16865120977}{1151056679735918592}$$

$$\langle \tau_6\tau_{10}\tau_{20}\rangle = \frac{201435244276049 69}{51061001021404654071038607360}$$

$$\langle \tau_6\tau_{11}\tau_{19}\rangle = \frac{17954012096899917 1}{25530500510702327035519303680 0}$$

$$\langle \tau_6\tau_{12}\tau_{15}\rangle = \frac{645431327778043}{17152482364014743204659200}$$

$$\langle \tau_6\tau_{13}\tau_{14}\rangle = \frac{370525355790311}{85762411820073716023296 00}$$

$$\langle \tau_6\tau_{14}\tau_{16}\rangle = \frac{16968031902890951}{93518316889019512950620160 00}$$

$$\langle {\tau_6\tau_{15}}^2\rangle = \frac{3284135262220031}{17003330343458093263749120 00}$$

$$\langle \tau_6\tau_{16}\tau_{17}\rangle = \frac{9241810139620751868061286 4000}$$

$$\langle \tau_6\tau_{17}\tau_{22}\rangle = \frac{18584828321010422089}{29964477810162373762029107281920000}$$

$$\langle \tau_6\tau_{19}\tau_{20}\rangle = \frac{38877451675127539 0013}{46744585383853303068765407359795200 00}$$

$$\langle {\tau_7}^3\rangle = \frac{8814691}{3210529996800}$$

$$\langle {\tau_7}^2\tau_{16}\rangle = \frac{91560179701}{32221598206604083 2000}$$

$$\langle \tau_7\tau_8\tau_9\rangle = \frac{1679060887}{6175225125273600}$$

$$\langle \tau_7\tau_8\tau_{18}\rangle = \frac{1517565628517}{11268353201395256524 8000}$$

$$\langle \tau_7\tau_9\tau_{14}\rangle = \frac{1028141503849}{11424021182341447 68000}$$

$$\langle \tau_7{\tau_{10}}^2\rangle = \frac{142227368603 3}{6330811738547552256 0}$$

$$\langle \tau_7\tau_{10}\tau_{13}\rangle = \frac{9938691090457}{7996814827639013376000}$$

$$\langle \tau_7\tau_{10}\tau_{16}\rangle = \frac{4980158843027}{1139355712585520381952 00}$$

$$\langle \tau_7\tau_{10}\tau_{19}\rangle = \frac{1376470684118437933}{12765250255351163511775965184000}$$

$$\langle \tau_7\tau_{10}\tau_{22}\rangle = \frac{11102336601883907}{55702910205168713532042117 1200}$$

$$\langle \tau_7\tau_{11}\tau_{12}\rangle = \frac{11667166895743}{7996814827639013376000}$$

$$\langle \tau_7\tau_{11}\tau_{15}\rangle = \frac{134464317788107}{2144060295501842900582400}$$

$$\langle \tau_7\tau_{11}\tau_{18}\rangle = \frac{179539670079186899}{9819423273347048859815116 8000}$$

$$\langle \tau_7\tau_{11}\tau_{21}\rangle = \frac{4344392757300223}{1114058204103374270640842342 40}$$

$$\langle \tau_7\tau_{12}\tau_{14}\rangle = \frac{476387883510529}{6125886558576694001664000}$$

$$\langle \tau_7\tau_{12}\tau_{17}\rangle = \frac{885729055722730831}{3273141091115682953271705600 00}$$

$$\langle \tau_7\tau_{12}\tau_{20}\rangle = \frac{5977884495667313}{8912465632826994165126738739 2}$$

$$\langle \tau_7{\tau_{13}}^2\rangle = \frac{108537290563007}{1299430482122329030656000}$$

$$\langle \tau_7\tau_{13}\tau_{16}\rangle = \frac{32804778301136311}{9351831688901951295062016000}$$

$$\langle \tau_7\tau_{13}\tau_{19}\rangle = \frac{20630745781051999}{2025560371097044128435895168 00}$$

$$\langle \tau_7\tau_{13}\tau_{22}\rangle = \frac{146630065607595893}{6698912076961466202546241536 0000}$$

$$\langle \tau_7\tau_{14}\tau_{15}\rangle = \frac{3393597870673783}{8501665171729046631874560 00}$$

$$\langle \tau_7\tau_{14}\tau_{18}\rangle = \frac{23476366133631989}{1713935698620575800985911296 00}$$

$$\langle \tau_7\tau_{14}\tau_{21}\rangle = \frac{35631106080908900609}{1049049631252165607318741424537 6000}$$

$$\langle \tau_7\tau_{15}\tau_{17}\rangle = \frac{934005965610353}{5713118995401919336619704320}$$

$$\langle \tau_7\tau_{15}\tau_{20}\rangle = \frac{494237923416371392 27}{1049049631252165607318741424537 6000}$$

$$\langle \tau_7{\tau_{16}}^2\rangle = \frac{38916914929495909}{2244439605336468310814883840 00}$$

$$\langle \tau_7\tau_{16}\tau_{19}\rangle = \frac{225152831249162739613}{3846515314591273893502051889971 2000}$$

$$\langle \tau_7\tau_{16}\tau_{22}\rangle = \frac{123898758494378164 9}{8561279374332106789151173509120000}$$

$$\langle \tau_7\tau_{17}\tau_{18}\rangle = \frac{2026375484920226665207}{3106800831016028913982426526515200 00}$$

$$\langle \tau_7\tau_{17}\tau_{21}\rangle = \frac{111508888270481616331}{5992895562032474752405821456384 0000}$$

$$\langle \tau_7\tau_{18}\tau_{20}\rangle = \frac{77754842538883236809}{3595737337219484851443492873830 40000}$$

$$\langle \tau_7{\tau_{19}}^2\rangle = \frac{10626495142470282604 93}{4674458538385330306876540735979520000}$$

$$\langle \tau_7\tau_{19}\tau_{22}\rangle = \frac{115935161364099220291}{1809467821310450441371564155863040 0000}$$

$$\langle \tau_7\tau_{20}\tau_{21}\rangle = \frac{254491817653934718 29}{3618935642620900882743128311726080000}$$

$$\langle {\tau_7\tau_{22}}^2\rangle = \frac{155861842966345884 19}{7632664264436809134512779712004096 0000}$$

$$\langle {\tau_8}^3\rangle = \frac{104256173}{343068062515200}$$

$$\langle {\tau_8}^2\tau_{11}\rangle = \frac{606258509}{26444493477642240}$$

$$\langle {\tau_8}^2\tau_{14}\rangle = \frac{63838947191}{63466784346341376000}$$

$$\langle {\tau_8}^2\tau_{17}\rangle = \frac{642238455259}{2191068678049077657 6000}$$

$$\langle {\tau_8}^2\tau_{20}\rangle = \frac{32914193308188 1}{5332184734900235387535360 00}$$

$$\langle \tau_8\tau_9\tau_{10}\rangle = \frac{50198200229}{180880335387079216}$$

$$\langle \tau_8\tau_9\tau_{13}\rangle = \frac{1753891748773}{1142402118234144768000}$$

$$\langle \tau_8\tau_9\tau_{16}\rangle = \frac{7909675910923}{1464885916181383348224 00}$$

$$\langle \tau_8\tau_9\tau_{19}\rangle = \frac{242906792635722497}{1823607179335588052537093120 00}$$

$$\langle \tau_8\tau_9\tau_{22}\rangle = \frac{4609968982291}{1872366729585550297586696192}$$

$$\langle \tau_8\tau_{10}\tau_{12}\rangle = \frac{15785016697657}{7996814827639013376000}$$

$$\langle \tau_8\tau_{10}\tau_{15}\rangle = \frac{79096736774 59}{9322001284790621306880 0}$$

$$\langle \tau_8\tau_{10}\tau_{18}\rangle = \frac{242906746815723701}{9819423273347048859815116 8000}$$

$$\langle \tau_8\tau_{10}\tau_{21}\rangle = \frac{2938854918876514 1}{5570291020516871353204211712 00}$$

$$\langle \tau_8{\tau_{11}}^2\rangle = \frac{9532018147}{4442674904243896320}$$

$$\langle \tau_8\tau_{11}\tau_{14}\rangle = \frac{40866651473243}{3573433825836404834304 00}$$

$$\langle \tau_8\tau_{11}\tau_{17}\rangle = \frac{21709058749847 33}{5455235151858947158878617600}$$

$$\langle \tau_8\tau_{11}\tau_{20}\rangle = \frac{3052433967649447}{3094606122509372974002339840 0}$$

$$\langle \tau_8\tau_{12}\tau_{13}\rangle = \frac{5688637764858331}{4288120591003685801164 8000}$$

$$\langle \tau_8\tau_{12}\tau_{16}\rangle = \frac{260508707218567867}{4675915844450975647531008000 0}$$

$$\langle \tau_8\tau_{12}\tau_{19}\rangle = \frac{3604314057316984 01}{2228116408206748541281684684800}$$

$$\langle \tau_8\tau_{12}\tau_{22}\rangle = \frac{10689334359670759224 7}{3074800643325312986987248650240000}$$

$$\langle \tau_8\tau_{13}\tau_{15}\rangle = \frac{2894541125666522 3}{4250832585864523315937280000}$$

$$\langle \tau_8\tau_{13}\tau_{18}\rangle = \frac{4004793344089115 3}{1713935698620575800985911296 00}$$

$$\langle \tau_8\tau_{13}\tau_{21}\rangle = \frac{356311142903520261 29}{6149601286650625973937449730048000}$$

$$\langle \tau_8{\tau_{14}}^2\rangle = \frac{1718980391533519}{2361573658813624064409600 00}$$

$$\langle \tau_8\tau_{14}\tau_{17}\rangle = \frac{7096616611798157}{2380466248084133056924876800 0}$$

$$\langle \tau_8\tau_{14}\tau_{20}\rangle = \frac{2553563199213949153 93}{2972307288547802554069767 3695232000}$$

$$\langle \tau_8\tau_{15}\tau_{16}\rangle = \frac{6867693483979253}{2040399641214971191649894 4000}$$

$$\langle \tau_8\tau_{15}\tau_{19}\rangle = \frac{4052751916718669696 23}{3566768746257363064883720843427 8400}$$

$$\langle \tau_8\tau_{15}\tau_{22}\rangle = \frac{530994759526434383}{1890152589138257343059349995520000}$$

$$\langle \tau_8\tau_{16}\tau_{18}\rangle = \frac{2026375956352372313671}{1509017546493499758220035741450240 00}$$

$$\langle \tau_8\tau_{16}\tau_{21}\rangle = \frac{1115088994374225187}{2910834987272916308311398993100 8000}$$

$$\langle \tau_8{\tau_{17}}^2\rangle = \frac{208266417926756253196 9}{1467100392424235876047256970854400 00}$$

$$\langle \tau_8\tau_{17}\tau_{20}\rangle = \frac{799147112937124431 29}{1697987075909201179848316079308800 00}$$

$$\langle \tau_8\tau_{18}\tau_{19}\rangle = \frac{3187949023152143633137}{6112753473273124247453937885511680000}$$

$$\langle \tau_8\tau_{18}\tau_{22}\rangle = \frac{347805518841347541563}{2366227150944435192562814665359360 0000}$$

$$\langle \tau_8\tau_{19}\tau_{21}\rangle = \frac{104341655647591540 1149}{6152190592455531500663318129934336 0000}$$

$$\langle {\tau_8\tau_{20}}^2\rangle = \frac{2026509120628706063 3}{1139294554158431759382095949987840000}$$

$$\langle \tau_8\tau_{21}\tau_{22}\rangle = \frac{1052067511573613619707}{1946329387431386329300758826561044480 0000}$$

$$\langle {\tau_9}^3\rangle = \frac{3658179233}{1192617595958722 56}$$

$$\langle {\tau_9}^2\tau_{12}\rangle = \frac{14954247302749}{6854412709404868608000}$$

$$\langle {\tau_9}^2\tau_{15}\rangle = \frac{17631482371}{1880067486008192 48}$$

$$\langle {\tau_9}^2\tau_{18}\rangle = \frac{1211169857514584 3}{4429815010532503245029376000}$$

$$\langle {\tau_9}^2\tau_{21}\rangle = \frac{3977398557196091}{6820764514918617983515 3612800}$$

$$\langle \tau_9\tau_{10}\tau_{11}\rangle = \frac{623093076061}{2399044448291704012 80}$$

$$\langle \tau_9\tau_{10}\tau_{14}\rangle = \frac{2277398902271}{1645059050257168465920 0}$$

$$\langle \tau_9\tau_{10}\tau_{17}\rangle = \frac{2489626434197983}{5168117512287920452534272 00}$$

$$\langle \tau_9\tau_{10}\tau_{20}\rangle = \frac{1995320626219089 47}{1671087306155061405961263513600}$$

$$\langle \tau_9\tau_{11}\tau_{13}\rangle = \frac{6604457586827}{3783635815591487471616 00}$$

$$\langle \tau_9\tau_{11}\tau_{16}\rangle = \frac{541223173865389}{7383025017554172075048960 0}$$

$$\langle \tau_9\tau_{11}\tau_{19}\rangle = \frac{3556889280049257 17}{1671087306155061405961263513600}$$

$$\langle \tau_9\tau_{11}\tau_{22}\rangle = \frac{166777644458165 59}{3646008667974679431188211302400}$$

$$\langle {\tau_9\tau_{12}}^2\rangle = \frac{2853126031689779}{1513454326236594988646400 0}$$

$$\langle \tau_9\tau_{12}\tau_{15}\rangle = \frac{1298935659881959 7}{1342368185009849468190720000}$$

$$\langle \tau_9\tau_{12}\tau_{18}\rangle = \frac{6829227504898863 7}{2056722838344690961183093555 20}$$

$$\langle \tau_9\tau_{12}\tau_{21}\rangle = \frac{1320878952321113719 9}{1604243813908858949722812973056000}$$

$$\langle \tau_9\tau_{13}\tau_{14}\rangle = \frac{7456852567799473}{6711840925049247340953600000}$$

$$\langle \tau_9\tau_{13}\tau_{17}\rangle = \frac{1949702265597337 21}{4284839246551439502464778240 00}$$

$$\langle \tau_9\tau_{13}\tau_{20}\rangle = \frac{1051811009829490905 7}{8021219069544294748614064865280 00}$$

$$\langle \tau_9\tau_{14}\tau_{16}\rangle = \frac{336155556461287161}{6121198923644913574946832000}$$

$$\langle \tau_9\tau_{14}\tau_{19}\rangle = \frac{4312425151948762556 09}{2326153530167845477098078810931200 0}$$

$$\langle \tau_9\tau_{14}\tau_{22}\rangle = \frac{1115089054007536140 1}{2432809300213434451228002091008000 0}$$

$$\langle {\tau_9\tau_{15}}^2\rangle = \frac{1301247354919361}{2225890517689059481799884800}$$

$$\langle \tau_9\tau_{15}\tau_{18}\rangle = \frac{8346629338837553442 1}{3578697738719762272458582786048000}$$

$$\langle \tau_9\tau_{15}\tau_{21}\rangle = \frac{8028641193016057777}{1206673412905863487809089037139968 0}$$

$$\langle \tau_9\tau_{16}\tau_{17}\rangle = \frac{5046164786816404429}{1929689957152812990051196600320 00}$$

$$\langle \tau_9\tau_{16}\tau_{20}\rangle = \frac{2876929758745387706 59}{3318351885491124591474994852134912 00}$$

$$\langle \tau_9\tau_{17}\tau_{19}\rangle = \frac{5897706006680756183911 9}{5807115799609468035081240991236096 0000}$$

$$\langle \tau_9\tau_{17}\tau_{22}\rangle = \frac{6434402331385664060573}{2247915793397213432934673932091392 00000}$$

$$\langle {\tau_9\tau_{18}}^2\rangle = \frac{5738308548849267266587}{5360414584254893570844222453448704 0000}$$

$$\langle \tau_9\tau_{18}\tau_{21}\rangle = \frac{1878149869841034769487}{5394997904153312239043217437019340 80000}$$

$$\langle \tau_9\tau_{19}\tau_{20}\rangle = \frac{1346007406590511332909 61}{3506748637699652955378091334062571520000}$$

$$\langle \tau_9\tau_{20}\tau_{22}\rangle = \frac{4679886634808379203905 9}{3825543968399621405867008728068259840 0000}$$

$$\langle {\tau_9\tau_{21}}^2\rangle = \frac{1959022312278185644 35 9}{1530217587359848562346803491227303936 0000}$$

$$\langle {\tau_{10}}^{3} \rangle = \frac{682434427273}{2399044448291704012 80}$$

$$\langle {\tau_{10}}^{2}\tau_{13} \rangle = \frac{293531282249}{1535388446906690568192}$$

$$\langle {\tau_{10}}^{2}\tau_{16} \rangle = \frac{4149376522392359}{5168117512287920452534272 00}$$

$$\langle {\tau_{10}}^{2}\tau_{19} \rangle = \frac{272694805367417681 9}{1169761114308542984172884459520 0}$$

$$\langle {\tau_{10}}^{2}\tau_{22} \rangle = \frac{156276817283671919}{3119362971489447957794358558720 0}$$

$$\langle \tau_{10}\tau_{11}\tau_{12} \rangle = \frac{594401002226287}{2648545070914041230131200}$$

$$\langle \tau_{10}\tau_{11}\tau_{15} \rangle = \frac{541223073795413}{4698288647534473138667520 00}$$

$$\langle \tau_{10}\tau_{11}\tau_{18} \rangle = \frac{355688890808287217}{8998162417758022955176034304 00}$$

$$\langle \tau_{10}\tau_{11}\tau_{21} \rangle = \frac{550366193566912447}{5614853348681006324029845405696 00}$$

$$\langle \tau_{10}\tau_{12}\tau_{14} \rangle = \frac{67111662223673413}{4698288647534473138667520 000}$$

$$\langle \tau_{10}\tau_{12}\tau_{17} \rangle = \frac{1754731878725777269}{2999387472586007651725344768 000}$$

$$\langle \tau_{10}\tau_{12}\tau_{20} \rangle = \frac{946629858263333583 73}{5614853348681006324029845405696 000}$$

$$\langle {\tau_{10}}^{2}\tau_{13} \rangle = \frac{72082892376709001}{4698288647534473138667520 000}$$

$$\langle \tau_{10}\tau_{13}\tau_{16} \rangle = \frac{324950341754567773}{4284839246551439502464778240 00}$$

$$\langle \tau_{10}\tau_{13}\tau_{19} \rangle = \frac{1774660442594675773}{6931917714420995461765241241 6000}$$

$$\langle {\tau_{10}}^{2}\tau_{22} \rangle = \frac{567325988661914464 7}{8962981632365284820313691914240 000}$$

$$\langle \tau_{10}\tau_{14}\tau_{15} \rangle = \frac{6723110603201053}{7790616811911708186299596800}$$

$$\langle \tau_{10}\tau_{14}\tau_{18} \rangle = \frac{8624849786043381007 3}{2505088417103833590721007950233600}$$

$$\langle \tau_{10}\tau_{14}\tau_{21} \rangle = \frac{191158116689929973}{1946247440170747560982401672806 40}$$

$$\langle \tau_{10}\tau_{15}\tau_{17} \rangle = \frac{85784796310141999717}{2087573680919861325600839958528000}$$

$$\langle \tau_{10}\tau_{15}\tau_{20} \rangle = \frac{2876929657209861947 63}{2111678472585261103665905814994944 00}$$

$$\langle {\tau_{10}}^{2}\tau_{16} \rangle = \frac{9531644007720483731}{2186981951439854722058022813696 00}$$

$$\langle \tau_{10}\tau_{16}\tau_{19} \rangle = \frac{393180386136275725787}{2322846319843787214032496396494438400}$$

$$\langle \tau_{10}\tau_{16}\tau_{22} \rangle = \frac{2144800729358927523073}{4495831586794426865869347864182784 0000}$$

$$\langle \tau_{10}\tau_{17}\tau_{18} \rangle = \frac{589770579580012833835 39}{3126908507482021249659129764511744 00000}$$

$$\langle \tau_{10}\tau_{17}\tau_{21} \rangle = \frac{193032065680395182553 67}{3147082110756098806108543504927948800 00}$$

$$\langle \tau_{10}\tau_{18}\tau_{20} \rangle = \frac{1346007376939539161276 53}{1888249266453659283665126102956769280000}$$

$$\langle {\tau_{10}}^{2}\tau_{19} \rangle = \frac{1839543414882081246335461}{2454724046389757068764663933843800064 0000}$$

$$\langle \tau_{10}\tau_{19}\tau_{22} \rangle = \frac{213194832590994256155809}{8926269259599116613689687032159272960600 0}$$

$$\langle \tau_{10}\tau_{20}\tau_{21} \rangle = \frac{140396597076859672559273}{5355761555759469968213812219295563776 0000}$$

$$\langle {\tau_{10}}^{2}\tau_{22} \rangle = \frac{589403185632472504124012023}{6896078579195893531072104613564967917977600 0000}$$

$$\langle {\tau_{11}}^{3} \rangle = \frac{1076813553581}{4414241784856735383552}$$

$$\langle {\tau_{11}}^{2}\tau_{14} \rangle = \frac{121579105281169}{7830481079224121897779200}$$

$$\langle {\tau_{11}}^{2}\tau_{17} \rangle = \frac{353206909537043}{5554421245529643799491379 20}$$

$$\langle {\tau_{11}}^{2}\tau_{20} \rangle = \frac{171490918700645549}{9358088914468343873383075676160}$$

$$\langle \tau_{11}\tau_{12}\tau_{13} \rangle = \frac{846190571741320 87}{4698288647534473138667520000}$$

$$\langle \tau_{11}\tau_{12}\tau_{16} \rangle = \frac{381463464984181087}{4284839246551439502464778240 00}$$

$$\langle \tau_{11}\tau_{12}\tau_{19} \rangle = \frac{153406422592349699 27}{5104412135164551203663495823360 00}$$

$$\langle \tau_{11}\tau_{12}\tau_{22} \rangle = \frac{744343315348051147 1}{1001745005970237715211530272768000 0}$$

$$\langle \tau_{11}\tau_{13}\tau_{15} \rangle = \frac{8476965794180407}{7790616811911708186299596800}$$

$$\langle \tau_{11}\tau_{13}\tau_{18} \rangle = \frac{3749934749703609791}{8638235921047702036968992938400}$$

$$\langle \tau_{11}\tau_{13}\tau_{21} \rangle = \frac{168717817554990397 7}{1362373208119523292687681170964480}$$

$$\langle {\tau_{11}}^{2}\tau_{14} \rangle = \frac{1510264030382663}{1298436135318618031049932800}$$

$$\langle \tau_{11}\tau_{14}\tau_{17} \rangle = \frac{14014907694720347}{2530392340508922818910109040 64}$$

$$\langle \tau_{11}\tau_{14}\tau_{20} \rangle = \frac{833892662710240 27}{4541244027065077642292270569881 6}$$

$$\langle {\tau_{11}}^{2}\tau_{15} \rangle = \frac{3729773820791801459}{5964496231199603787430971310080 0}$$

$$\langle \tau_{11}\tau_{15}\tau_{19} \rangle = \frac{5128439876666195233901}{2111678472585261103665905814994944 00}$$

$$\langle \tau_{11}\tau_{15}\tau_{22} \rangle = \frac{279756618625461692339}{4087119624358569878063043512893440000}$$

$$\langle {\tau_{11}}^{2}\tau_{16} \rangle = \frac{5128439874285604478 41}{1786804861418297856948074151149568 00}$$

$$\langle \tau_{11}\tau_{16}\tau_{21} \rangle = \frac{119895693664549766231}{1284523310512693390248385104052224000}$$

$$\langle {\tau_{11}}^{2}\tau_{17} \rangle = \frac{263544826957789676692 1}{8685856965227836804068693790310400 00}$$

$$\langle \tau_{11}\tau_{17}\tau_{20} \rangle = \frac{200492243324220055758 7}{1748378950420054892282524169404416000 0}$$

$$\langle \tau_{11}\tau_{18}\tau_{19} \rangle = \frac{239904046869402299305323}{1888249266453659283665126102956769280000}$$

$$\langle {\tau_{11}}^{2}\tau_{18} \rangle = \frac{834240652223110680837 61}{2059908290676719218543773930498293760000 0}$$

$$\langle \tau_{11}\tau_{18}\tau_{22} \rangle = \frac{230636029653876267345146527}{1379215715839178706214420922712993583595520000}$$

$$\langle \tau_{11}\tau_{19}\tau_{21} \rangle = \frac{250272195661730101436783}{5355761555759469968213812219295563776 0000}$$

$$\langle {\tau_{11}}^{2}\tau_{19} \rangle = \frac{161997183294125992723}{3455294368419080814787597172736000}$$

$$\langle {\tau_{11}}^{2}\tau_{20} \rangle = \frac{430063937968879 19}{2210959363545634418196480000}$$

$$\langle {\tau_{20}}^{2}\tau_{21} \rangle = \frac{4374676590909577402108 9}{8926269259599116613689687032159272960000}$$

$$\langle \tau_{11}\tau_{20}\tau_{21} \rangle = \frac{184996782002648132 59}{2879411973682567345656333097728000}$$

$$\langle {\tau_{12}}^{3} \rangle = \frac{3854788756050049}{3280259710278613973178777600}$$

$$\langle {\tau_{12}}^{2}\tau_{15} \rangle = \frac{1106467123795507 3}{8200649275696534932946944000}$$

$$\langle {\tau_{12}}^{2}\tau_{18} \rangle = \frac{178380517959782424 73}{5923361774432709968207309438976000}$$

$$\langle {\tau_{12}}^{2}\tau_{21} \rangle = \frac{932047284978551507}{6968660911097305844949777581056000}$$

$$\langle \tau_{12}\tau_{13}\tau_{14} \rangle = \frac{924983912099163543959}{1192899246239920757486194262016000}$$

$$\langle \tau_{12}\tau_{13}\tau_{17} \rangle = \frac{107028310842041495531}{2824987923190984754068101424742400 00}$$

$$\langle \tau_{12}\tau_{13}\tau_{20} \rangle = \frac{126171585708213999 01}{5923361774432709968207309438976000}$$

$$\langle \tau_{12}\tau_{14}\tau_{16} \rangle = \frac{179029144720911675 1}{2168907720436219559006580774912000}$$

$$\langle \tau_{12}\tau_{14}\tau_{19} \rangle = \frac{485224925873832023 89}{3022407789441631506466788480122880 00}$$

$$\langle \tau_{12}\tau_{14}\tau_{22} \rangle = \frac{603301230770664299 81}{7108034129319251961848771326771200 00}$$

$$\langle \tau_{12}\tau_{15}\tau_{17} \rangle = \frac{110001319390902788419}{2589572262925069357895759639347200 0}$$

$$\langle \tau_{12}\tau_{15}\tau_{18} \rangle = \frac{135495782032970057189 09}{6835291462275689714624890870431744 0000}$$

$$\langle \tau_{12}\tau_{15}\tau_{21} \rangle = \frac{379152500337950077}{3077360658340570261129820997943296}$$

$$\langle \tau_{12}\tau_{16}\tau_{17} \rangle = \frac{823117444475791490208 49}{1373272193784479479029182620332195840000 0}$$

$$\langle \tau_{12}\tau_{16}\tau_{20} \rangle = \frac{273582049168650643884819157 }{9511832523028818663547730501468921266176000 00}$$

$$\langle \tau_{12}\tau_{17}\tau_{19} \rangle = \frac{236741241356908724769691}{1258832844302439522443147401971179520000}$$

$$\langle \tau_{12}\tau_{17}\tau_{22} \rangle = \frac{172187270815824971681597 3}{2142304622303787987285524887718225510400 00}$$

$$\langle \tau_{12}\tau_{18}\tau_{21} \rangle = \frac{126453320065328661434 09}{1734659613201447762984230678314352640000}$$

$$\langle \tau_{12}\tau_{19}\tau_{20} \rangle = \frac{149037748100504378055 1}{1634847849743427951225217405157376000 0}$$

$$\langle {\tau_{12}}^{2}\tau_{21} \rangle = \frac{114522718257345774380202557}{3804733009211527465419092200587568506470400 00}$$

$$\langle {\tau_{13}}^{3} \rangle = \frac{11884276233822961}{8200649275696534932946944000}$$

$$\langle {\tau_{13}}^{2}\tau_{16} \rangle = \frac{126883937204803861}{1523498398773845156431921152000}$$

$$\langle {\tau_{13}}^{2}\tau_{19} \rangle = \frac{191593888867594533601}{5923361774432709968207309438976000}$$

$$\langle {\tau_{13}}^{2}\tau_{22} \rangle = \frac{272077025135568828 61}{1923350411462856413206138123714560 00}$$

$$\langle \tau_{13}\tau_{14}\tau_{15} \rangle = \frac{102775989655148012 83}{1084453860218109779532903874560 00}$$

$$\langle \tau_{13}\tau_{14}\tau_{18} \rangle = \frac{198200574836901900 1}{4556432134179007667851776491520 00}$$

$$\langle \tau_{13}\tau_{14}\tau_{21} \rangle = \frac{237061461977698001 9}{3693388947035813368422513770496000}$$

$$\langle \tau_{13}\tau_{15}\tau_{17} \rangle = \frac{122223687853135319 27}{2354156602659153961723417853952000}$$

$$\langle \tau_{13}\tau_{15}\tau_{20} \rangle = \frac{128315035748946565187}{6539391398973711804900869620629504 00}$$

$$\langle \tau_{13}\tau_{16}\tau_{19} \rangle = \frac{125259158041366057301}{5138093242050773560993540416208896000}$$

$$\langle \tau_{13}\tau_{16}\tau_{22} \rangle = \frac{390843990298092670991}{5030301076133624465308361246638080 0000}$$

$$\langle \tau_{13}\tau_{17}\tau_{18} \rangle = \frac{138445170133014493356 1}{5096489248188014260904523894620160000}$$

$$\langle \tau_{13}\tau_{17}\tau_{21} \rangle = \frac{137186240605506306965371}{1373272193784479479029182620332195840000}$$

$$\langle \tau_{13}\tau_{18}\tau_{20} \rangle = \frac{956595948017276411219689}{8239633162706876874175095721993175040000 0}$$

$$\langle {\tau_{13}}^{2}\tau_{20} \rangle = \frac{435782598517426533081031 1}{3570507703839646645475874812863709184000 00}$$

$$\langle \tau_{13}\tau_{19}\tau_{22} \rangle = \frac{415439407749430746222023501}{9511832523028818663547730501468921266176000 00}$$

$$\langle \tau_{13}\tau_{20}\tau_{21} \rangle = \frac{911940163363683182524228 19}{1902366504605763732709546100293784253235200 00}$$

$$\langle {\tau_{13}}^{2}\tau_{22} \rangle = \frac{509990182123392200948579}{2924571462157770066984096741128247705600000 0}$$

$$\langle {\tau_{14}}^{3} \rangle = \frac{1831066487060472973}{1807423100363516299221506476000}$$

$$\langle {\tau_{14}}^2 \tau_{17} \rangle = \frac{23414499614949743}{42189186427583404331960893440000}$$

$$\langle \tau_{14} \tau_{15} \tau_{19} \rangle = \frac{3672285682784833909}{13237634410877959119232529596416000}$$

$$\langle \tau_{14} \tau_{16} \tau_{21} \rangle = \frac{4730560022267075714303}{39236348393842270829405217723777024000}$$

$$\langle \tau_{14} \tau_{18} \tau_{19} \rangle = \frac{1352428754551622110128037}{8239633162706876874175095721993175040000}$$

$$\langle \tau_{14} {\tau_{20}}^2 \rangle = \frac{32194931881766075533696 49}{45294440585851517445465383340328196505600000}$$

$$\langle {\tau_{15}}^2 \tau_{18} \rangle = \frac{218085649505684627}{624882121258835337305386490265600}$$

$$\langle \tau_{15} \tau_{16} \tau_{20} \rangle = \frac{1312348909533728275697}{78472696787684541658810435447554048000}$$

$$\langle \tau_{15} {\tau_{18}}^2 \rangle = \frac{523520808285285484759 57}{25352717423713467305154140683055923200 00}$$

$$\langle \tau_{15} \tau_{20} \tau_{22} \rangle = \frac{330823634008390084329784 7}{88140533031927277191175881094692706713600000}$$

$$\langle {\tau_{16}}^2 \tau_{19} \rangle = \frac{597847836467457614550 7}{28773322155484331941563826330769817600 0}$$

$$\langle \tau_{16} \tau_{17} \tau_{21} \rangle = \frac{3465846005232923308730179}{36254385915910258246370421176769970176000 00}$$

$$\langle \tau_{16} \tau_{19} \tau_{22} \rangle = \frac{9042512662545855075508619}{19390917267024000982058693840832395476992 00000}$$

$$\langle {\tau_{17}}^3 \rangle = \frac{2257180607411319191 29}{92222186395783115197319956188364800 00}$$

$$\langle \tau_{17} \tau_{18} \tau_{22} \rangle = \frac{1356376899409285039975854 59}{26103157859455385937386703247274378526720000 00}$$

$$\langle \tau_{17} \tau_{21} \tau_{22} \rangle = \frac{1057844571695605417974075443}{39833139715369994283918019457979768215961600000 0}$$

$$\langle \tau_{18} \tau_{19} \tau_{20} \rangle = \frac{1233649494878161646004587 93}{17704750548152348722749242202499143696384000 00}$$

$$\langle {\tau_{19}}^3 \rangle = \frac{2982901214510928468022863079}{40720926260750402062323257065748030501683200000}$$

$$\langle \tau_{19} {\tau_{22}}^2 \rangle = \frac{1364569522408068750074196767329}{86025591609299155265026619115085204013249986560000 00}$$

$$\langle {\tau_{14}}^2 \tau_{20} \rangle = \frac{462767707930924427}{22062724018129931865387543932736000}$$

$$\langle \tau_{14} \tau_{15} \tau_{22} \rangle = \frac{121296410843128764497}{13719002934909884905386439763558400000}$$

$$\langle \tau_{14} {\tau_{17}}^2 \rangle = \frac{9322505236576039747 97}{26898137698770075265884987221606400 00}$$

$$\langle \tau_{14} \tau_{18} \tau_{22} \rangle = \frac{4297649046556797010006113 7}{73167942484837066642674850011299394355200000}$$

$$\langle \tau_{14} \tau_{21} \tau_{22} \rangle = \frac{691652202242122316308685 93}{25560754579258910385441005517460884946944000 00}$$

$$\langle {\tau_{15}}^2 \tau_{21} \rangle = \frac{183118452519950336503}{14267763052306280301601897354100736000}$$

$$\langle \tau_{15} \tau_{17} \tau_{19} \rangle = \frac{53806305293331329572609}{27465443875689589580583652406644391680000}$$

$$\langle \tau_{15} \tau_{18} \tau_{21} \rangle = \frac{249540912396058411691509 39}{29267176993934826657069940004519757742080000}$$

$$\langle \tau_{15} {\tau_{21}}^2 \rangle = \frac{1384843119108209483392843}{35256213212770910876470352437877082685440000}$$

$$\langle {\tau_{16}}^2 \tau_{22} \rangle = \frac{3850940005682440457520 77}{51791979879871797494814887395385671680000 0}$$

$$\langle \tau_{16} \tau_{18} \tau_{20} \rangle = \frac{1277411813364854568122139 31}{11497819533331539043848905001775619112960000 0}$$

$$\langle \tau_{16} \tau_{20} \tau_{21} \rangle = \frac{4315090878224131969435 63}{84308335943582612965472581916662589030400000}$$

$$\langle {\tau_{17}}^2 \tau_{20} \rangle = \frac{7096732296628967355916 13}{60423976526517097077284035294616616960000 0}$$

$$\langle \tau_{17} \tau_{19} \tau_{21} \rangle = \frac{1769187260091150090619892 3}{29507917580253914537915403670831906160640000 0}$$

$$\langle {\tau_{18}}^3 \rangle = \frac{2038334958495643903899304 67}{14858720627689898918204738771525415469056000 0}$$

$$\langle \tau_{18} {\tau_{20}}^2 \rangle = \frac{2543559175190251619286112 93}{82413392514558608863278660947544348033024000000}$$

$$\langle {\tau_{19}}^2 \tau_{22} \rangle = \frac{2826176861310661976935026 47}{87103585584492838635985576611225733693440000000}$$

$$\langle {\tau_{20}}^3 \rangle = \frac{1333817616256765359341572 63}{35712470089642063840754086410602550814310400000}$$

$$\langle \tau_{14} \tau_{15} \tau_{16} \rangle = \frac{2107304965306301063}{33630808609416485167477397913600 0}$$

$$\langle \tau_{14} \tau_{16} \tau_{18} \rangle = \frac{1851131762136775183 1}{56462563099459050120808136441856000}$$

$$\langle \tau_{14} \tau_{17} \tau_{20} \rangle = \frac{2607872832951318082813}{17606053766467685628579264363233280 0000}$$

$$\langle \tau_{14} \tau_{19} \tau_{21} \rangle = \frac{1289294713954774197790723 91}{19023665046057637327095461002937842532350 0000}$$

$$\langle {\tau_{15}}^3 \rangle = \frac{407865477597219179}{61146924744393609395413450752000}$$

$$\langle \tau_{15} \tau_{16} \tau_{17} \rangle = \frac{784502544481556653 7}{20048301390387633738547816562688000}$$

$$\langle \tau_{15} \tau_{17} \tau_{22} \rangle = \frac{4621128007278705212064 1}{65917065301655014993400765775945400320000}$$

$$\langle \tau_{15} \tau_{19} \tau_{20} \rangle = \frac{2554823626872077535258441 1}{27176664351510910467279230004196917903360000}$$

$$\langle {\tau_{16}}^3 \rangle = \frac{1188640218358884251}{28640430557696619626496880803840 00}$$

$$\langle \tau_{16} \tau_{17} \tau_{18} \rangle = \frac{5380630528745543461531 7}{23239990971737345029724628959467929600 00}$$

$$\langle \tau_{16} \tau_{19} {}^2 \rangle = \frac{1745796144893777253808092707}{14947165393331000757003576502308304846848000 00}$$

$$\langle \tau_{16} {\tau_{22}}^2 \rangle = \frac{3917942858102068055397487 1}{18968161769223806801865723551418937245696000000}$$

$$\langle \tau_{17} \tau_{18} \tau_{19} \rangle = \frac{1047477686982113541721818679}{80484736733320773306942335012429333790720000 0}$$

$$\langle \tau_{17} {\tau_{20}}^2 \rangle = \frac{4741805420689517701880449}{75409122705093337152450476047681537966080000}$$

$$\langle {\tau_{18}}^2 \tau_{21} \rangle = \frac{3959154192892009371187328 1}{62647578862692926249728087793458508464128000 00}$$

$$\langle \tau_{18} {\tau_{21}}^2 \rangle = \frac{6613327336916101181750 77}{20475377022250586052988487191936483983360000 0}$$

$$\langle \tau_{19} \tau_{20} \tau_{21} \rangle = \frac{7630677525554176931346921 71}{21427482053785238304452451846361530488586240000 0}$$

$$\langle \tau_{20} \tau_{21} \tau_{22} \rangle = \frac{5990793025211453654394615727 1}{34410236643719662106010647646034081605299946240000}$$

The nonzero four point intersection numbers $\langle \tau_{j_1}\tau_{j_2}\tau_{j_3}\tau_{j_4}\rangle$ with $2 \le j_1 \le j_2 \le j_3 \le j_4 \le 9$:

$$\langle {\tau_2}^3\tau_4\rangle = \frac{53}{1152}$$
$$\langle {\tau_2}^3\tau_7\rangle = \frac{4907}{1382400}$$
$$\langle {\tau_2}^2{\tau_3}^2\rangle = \frac{205}{3456}$$
$$\langle {\tau_2}^2\tau_3\tau_6\rangle = \frac{7897}{1036800}$$
$$\langle {\tau_2}^2\tau_3\tau_9\rangle = \frac{117871}{298598400}$$
$$\langle {\tau_2}^2\tau_4\tau_5\rangle = \frac{7597}{691200}$$
$$\langle {\tau_2}^2\tau_4\tau_8\rangle = \frac{27629}{33177600}$$
$$\langle {\tau_2}^2\tau_5\tau_7\rangle = \frac{14237}{11059200}$$
$$\langle {\tau_2}^2{\tau_6}^2\rangle = \frac{295693}{199065600}$$
$$\langle {\tau_2}^2\tau_6\tau_9\rangle = \frac{155893}{1433272320}$$
$$\langle {\tau_2}^2\tau_7\tau_8\rangle = \frac{365669}{2654208000}$$
$$\langle {\tau_2}^2{\tau_9}^2\rangle = \frac{606960439}{56757583872000}$$
$$\langle \tau_2{\tau_3}^2\tau_5\rangle = \frac{1837}{129600}$$
$$\langle \tau_2{\tau_3}^2\tau_8\rangle = \frac{747709}{696729600}$$
$$\langle \tau_2\tau_3{\tau_4}^2\rangle = \frac{177}{10240}$$
$$\langle \tau_2\tau_3\tau_4\tau_7\rangle = \frac{2821883}{1393459200}$$
$$\langle \tau_2\tau_3\tau_5\tau_6\rangle = \frac{1924831}{696729600}$$
$$\langle \tau_2\tau_3\tau_5\tau_9\rangle = \frac{5072357}{25082265600}$$
$$\langle \tau_2\tau_3\tau_6\tau_8\rangle = \frac{4942457}{16721510400}$$
$$\langle \tau_2\tau_3{\tau_7}^2\rangle = \frac{112020239}{334430208000}$$
$$\langle \tau_2\tau_3\tau_8\tau_9\rangle = \frac{3847708957}{1324343623680000}$$
$$\langle \tau_2{\tau_4}^2\tau_6\rangle = \frac{4699193}{1393459200}$$
$$\langle \tau_2{\tau_4}^2\tau_9\rangle = \frac{24779479}{100329062400}$$
$$\langle \tau_2\tau_4{\tau_5}^2\rangle = \frac{92599}{23224320}$$
$$\langle \tau_2\tau_4\tau_5\tau_8\rangle = \frac{264259}{619315200}$$
$$\langle \tau_2\tau_4\tau_6\tau_7\rangle = \frac{7464077}{13377208320}$$
$$\langle \tau_2\tau_4\tau_7\tau_9\rangle = \frac{440304889}{8026324992000}$$
$$\langle \tau_2\tau_4{\tau_8}^2\rangle = \frac{82019143}{1337720832000}$$
$$\langle \tau_2{\tau_5}^2\tau_7\rangle = \frac{408461}{619315200}$$
$$\langle \tau_2\tau_5{\tau_6}^2\rangle = \frac{254515}{334430208}$$
$$\langle \tau_2\tau_5\tau_6\tau_9\rangle = \frac{180147253}{2407897497600}$$
$$\langle \tau_2\tau_5\tau_7\tau_8\rangle = \frac{42255967}{445906944000}$$
$$\langle \tau_2\tau_5{\tau_9}^2\rangle = \frac{647709563}{68109100646400}$$
$$\langle \tau_2{\tau_6}^2\tau_8\rangle = \frac{175531261}{1605264998400}$$
$$\langle \tau_2\tau_6{\tau_7}^2\rangle = \frac{663092951}{5350883328000}$$
$$\langle \tau_2\tau_6\tau_8\tau_9\rangle = \frac{11474717}{825564856320}$$
$$\langle \tau_2{\tau_7}^2\tau_9\rangle = \frac{28609738103}{1816242683904000}$$
$$\langle \tau_2\tau_7{\tau_8}^2\rangle = \frac{1776434317}{100902371328000}$$
$$\langle \tau_2\tau_8{\tau_9}^2\rangle = \frac{29014006349}{13076947324108800}$$
$$\langle {\tau_3}^3\tau_4\rangle = \frac{185251}{8294400}$$
$$\langle {\tau_3}^3\tau_7\rangle = \frac{1455581}{557383680}$$
$$\langle {\tau_3}^2\tau_4\tau_6\rangle = \frac{807977}{185794560}$$
$$\langle {\tau_3}^2\tau_4\tau_9\rangle = \frac{446929871}{1404606873600}$$
$$\langle {\tau_3}^2{\tau_5}^2\rangle = \frac{1790449}{348364800}$$
$$\langle {\tau_3}^2\tau_5\tau_8\rangle = \frac{32168387}{8852528640 0}$$
$$\langle {\tau_3}^2{\tau_6}\tau_7\rangle = \frac{224368483}{312134860800}$$
$$\langle {\tau_3}^2\tau_7\tau_9\rangle = \frac{261919892621}{3708162146304000}$$
$$\langle {\tau_3}^2{\tau_8}^2\rangle = \frac{24394460551}{3090135121920 00}$$
$$\langle \tau_3{\tau_4}^2\tau_5\rangle = \frac{624443}{99532800}$$
$$\langle \tau_3{\tau_4}^2\tau_8\rangle = \frac{157148641}{234101145600}$$
$$\langle \tau_3\tau_4\tau_5\tau_7\rangle = \frac{242896603}{234101145600}$$
$$\langle \tau_3\tau_4{\tau_6}^2\rangle = \frac{6406985}{5350883328}$$
$$\langle \tau_3\tau_4\tau_6\tau_9\rangle = \frac{7934017549}{67421129932800}$$
$$\langle \tau_3\tau_4\tau_7\tau_8\rangle = \frac{930495989}{6242697216000}$$
$$\langle \tau_3\tau_4{\tau_9}^2\rangle = \frac{38028187861}{2542739757465600}$$
$$\langle \tau_3{\tau_5}^2\tau_6\rangle = \frac{2366359}{1672151040}$$
$$\langle \tau_3{\tau_5}^2\tau_9\rangle = \frac{1172122057}{8427641241600}$$
$$\langle \tau_3\tau_5\tau_6\tau_8\rangle = \frac{1142098963}{5618427494400}$$
$$\langle \tau_3\tau_5{\tau_7}^2\rangle = \frac{1438143943}{6242697216000}$$
$$\langle \tau_3\tau_5\tau_8\tau_9\rangle = \frac{4105508153}{158921234841600}$$
$$\langle \tau_3{\tau_6}^2\tau_7\rangle = \frac{11948459099}{44947419955200}$$
$$\langle \tau_3\tau_6\tau_7\tau_9\rangle = \frac{6872104145}{203419180597248}$$
$$\langle \tau_3\tau_6{\tau_8}^2\rangle = \frac{285736471}{7567677849600}$$
$$\langle \tau_3{\tau_7}^2\tau_8\rangle = \frac{2355115237}{550376570880 00}$$
$$\langle \tau_3{\tau_7}^2{\tau_9}^2\rangle = \frac{3290451525797}{610257541791744000}$$
$$\langle \tau_3{\tau_8}^2\tau_9\rangle = \frac{919391811931}{152564385447936000}$$
$$\langle {\tau_4}^4\rangle = \frac{84659}{11059200}$$
$$\langle {\tau_4}^3\tau_7\rangle = \frac{59322643}{46820229120}$$
$$\langle {\tau_4}^2\tau_5\tau_6\rangle = \frac{80918813}{46820229120}$$
$$\langle {\tau_4}^2\tau_5\tau_9\rangle = \frac{19481381}{114661785600}$$
$$\langle {\tau_4}^2{\tau_6}\tau_8\rangle = \frac{930136433}{37456183296 00}$$
$$\langle {\tau_4}^2{\tau_7}^2\rangle = \frac{5270462621}{18728091648000}$$
$$\langle {\tau_4}^2\tau_8\tau_9\rangle = \frac{1605173617}{50854795149312}$$
$$\langle {\tau_4}{\tau_5}^3\rangle = \frac{7970521}{3901685760}$$
$$\langle {\tau_4}{\tau_5}^2\tau_8\rangle = \frac{91611143}{312134860800}$$
$$\langle {\tau_4}\tau_5\tau_6\tau_7\rangle = \frac{1437601939}{37456183296 00}$$
$$\langle {\tau_4}\tau_5\tau_7\tau_9\rangle = \frac{12404250983}{2542739757465 60}$$
$$\langle {\tau_4}\tau_5{\tau_8}^2\rangle = \frac{3851017033}{70631659929600}$$
$$\langle {\tau_4}{\tau_6}^3\rangle = \frac{3981318281}{8989483991040}$$
$$\langle {\tau_4}{\tau_6}^2\tau_9\rangle = \frac{247846157}{4403012567040}$$
$$\langle {\tau_4}\tau_6\tau_7\tau_8\rangle = \frac{1098763513}{15410543984640}$$
$$\langle {\tau_4}\tau_6{\tau_9}^2\rangle = \frac{3289825492741}{3661545250750 46400}$$
$$\langle {\tau_4}{\tau_7}^3\rangle = \frac{10376946599}{128421199872000}$$
$$\langle {\tau_4}\tau_7\tau_8\tau_9\rangle = \frac{3472558154917}{3051287708958 72000}$$
$$\langle {\tau_4}{\tau_8}^3\rangle = \frac{19601536903}{1541054398464000}$$
$$\langle {\tau_4}{\tau_9}^3\rangle = \frac{46389657079979}{2636312580540 3340800}$$
$$\langle {\tau_5}^3\tau_7\rangle = \frac{424777463}{936404582400}$$
$$\langle {\tau_5}^2{\tau_6}^2\rangle = \frac{588180611}{1123685498880}$$
$$\langle {\tau_5}^2\tau_6\tau_9\rangle = \frac{48057059}{722369249280}$$
$$\langle {\tau_5}^2\tau_7\tau_8\rangle = \frac{541079213}{6421059993600}$$
$$\langle {\tau_5}^2{\tau_9}^2\rangle = \frac{97201583659}{9153863126876160}$$
$$\langle {\tau_5}\tau_6{\tau_7}^2\rangle = \frac{8490865079}{77052719923200}$$
$$\langle {\tau_5}\tau_6\tau_8\tau_9\rangle = \frac{473551272401}{3051287708958 7200}$$
$$\langle {\tau_5}{\tau_7}^2\tau_9\rangle = \frac{5366854169191}{3051287708958 72000}$$
$$\langle {\tau_5}\tau_7{\tau_8}^2\rangle = \frac{12342107077}{627836977152000}$$
$$\langle {\tau_5}\tau_8{\tau_9}^2\rangle = \frac{667744953581}{2196927150450 27840}$$
$$\langle {\tau_6}^3\tau_7\rangle = \frac{7837998359}{61642175938560}$$
$$\langle {\tau_6}^2\tau_7\tau_9\rangle = \frac{1651374355313}{8136767223889 9200}$$
$$\langle {\tau_6}^2{\tau_8}^2\rangle = \frac{92282814913}{4068383611944960}$$
$$\langle {\tau_6}{\tau_7}^2\tau_8\rangle = \frac{475390979681}{1849265278156 8000}$$
$$\langle {\tau_6}\tau_7{\tau_9}^2\rangle = \frac{768425554378511}{1933295892396 24499200}$$
$$\langle {\tau_6}{\tau_8}^2\tau_9\rangle = \frac{35784554826989}{8055399551651 020800}$$
$$\langle {\tau_7}^4\rangle = \frac{538769781889}{1849265278156 8000}$$
$$\langle {\tau_7}^2\tau_8\tau_9\rangle = \frac{270370425938309}{5370266367767 3472000}$$
$$\langle \tau_7{\tau_8}^3\rangle = \frac{7194739537639}{1278634849468 416000}$$
$$\langle \tau_7{\tau_9}^3\rangle = \frac{1863628639016059}{1988532917893 280563200}$$
$$\langle {\tau_8}^2{\tau_9}^2\rangle = \frac{1577939527948 7}{1506464331737 3337600}$$

References

[1] Arbarello, E., Cornalba, M. (1996). Combinatorial and algebro-geometric cohomology classes on the moduli spaces of curves. Journal of Algebraic Geometry, **5** (4): 705–750.

[2] Bergere, M., Borot, G., Eynard, B. (2013, December). Rational differential systems, loop equations, and application to the q-th reductions of KP. In Annales Henri Poincaré (pp. 1-70). Springer Basel.

[3] Babelon, O., Bernard, D., Talon, M. (2003). Introduction to classical integrable systems. Cambridge University Press.

[4] Balogh F., Bertola, M., Bothner T., Yang, D. in preparation.

[5] Balogh, F., Yang, D. (2014). Geometric interpretation of Zhou's explicit formula for the Witten–Kontsevich tau-function. arXiv preprint arXiv: 1412.4419.

[6] Bergere, M., Eynard, B. (2009). Determinantal formulae and loop equations. arXiv preprint arXiv: 0901.3273.

[7] Bertola, M. (2010). The dependence on the monodromy data of the isomonodromic tau-function. Communications in Mathematical Physics, **294** (2): 539–579.

[8] Bertola, M., Dubrovin, B., Yang, D. in preparation.

[9] Bertola, M., Yang, D. (2015). The partition function of the extended r-reduced Kadomtsev–Petviashvili hierarchy. J. Phys. A: Math. Theor. **48** (2015) 195205.

[10] Brézin, E., Hikami, S. (2007). Vertices from replica in a random matrix theory. Journal of Physics A: Mathematical and Theoretical, **40** (45): 13545–13566.

[11] Brézin, E., Hikami, S. (2015). Random Matrix, Singularities and Open/Close Intersection Numbers. arXiv preprint arXiv: arXiv:1502.01416.

[12] Buryak, A. (2014). Open intersection numbers and the wave function of the KdV hierarchy. arXiv preprint arXiv: 1409.7957.

[13] Buryak, A. (2013). Dubrovin–Zhang hierarchy for the Hodge integrals. arXiv preprint arXiv: 1308.5716.

[14] Carlet, G., Dubrovin, B., Zhang, Y. (2004). The extended Toda hierarchy. Mosc. Math. J, **4** (2): 313–332.

[15] Date, E., Jimbo, M., Kashiwara, M., Miwa, T. (1981). Transformation groups for soliton equations, Proceedings of RIMS.

[16] Dickey, L. A. (2003). Soliton Equations and Hamiltonian Systems (Vol. 26). 2nd edition. World Scientific.

[17] Dijkgraaf, R., Verlinde, H., Verlinde, E. (1991). Loop equations and Virasoro constraints in non-perturbative two-dimensional quantum gravity. Nuclear Physics B, **348** (3), 435–456.

[18] Dubrovin, B. (1975). Periodic problems for the Korteweg–de Vries equation in the class of finite band potentials. Functional Analysis and its Applications, **9** (3): 215–223.

[19] Dubrovin, B. (1996). Geometry of 2D topological field theories. In "Integrable Systems and Quantum Groups", Editors: Francaviglia, M., Greco, S. Springer Lecture Notes in Math. **1620**: 120–348.

[20] Dubrovin, B. (2013). Gromov–Witten invariants and integrable hierarchies of topological type. In Topology, Geometry, Integrable Systems, and Mathematical Physics: Novikov's Seminar 2012–2014 (Vol. 234, p. 141). American Mathematical Soc..

[21] Dubrovin, B., Liu, S.-Q., Yang, D., Zhang, Y. (2014). Hodge integrals and tau-symmetric integrable hierarchies of Hamiltonian evolutionary PDEs. arXiv preprint arXiv:1409.4616.

[22] Dubrovin, B., Zhang, Y. (2001). Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov-Witten invariants. arXiv preprint arXiv:math/0108160.

[23] Eynard, B., Orantin, N. (2007). Invariants of algebraic curves and topological expansion. Communications in Number Theory and Physics, **1** (2007) 347-452.

[24] Eynard, B., Orantin, N. (2007). Weil–Petersson volume of moduli spaces, Mirzakhani's recursion and matrix models. arXiv preprint arXiv:0705.3600.

[25] Faber, C. (1999). A conjectural description of the tautological ring of the moduli space of curves. In Moduli of curves and abelian varieties (pp. 109–129). Vieweg + Teubner Verlag.

[26] Givental, A. (2003). A_n-singularities and nKdV hierarchies. Moscow Mathematical Journal, **3** (2): 475–505.

[27] Hirota, R. (1971). Exact solution of the Korteweg–de Vries equation for multiple collisions of solitons. Physical Review Letters, **27** (18): 1192–1194.

[28] Itzykson, C., Zuber, J.-B. (1992). Combinatorics of the modular group. II. The Kontsevich integrals. Internat. J. Modern Phys. A, **7** (23): 5661–5705.

[29] Jimbo, M., Miwa, T., Ueno K. (1981). Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. I. General theory and τ-function. Phys. D, **2** (2): 306–352.

[30] Kabanov, A., Kimura, T. (1998). Intersection numbers and rank one cohomological field theories in genus one. Comm. Math. Phys. **194** (3): 651–674.

[31] Kac, V., Schwarz, A. (1991). Geometric interpretation of the partition function of 2D gravity. Physics Letters B, **257** (3): 329–334.

[32] Kaufmann, R., Manin, Y., Zagier, D. (1996). Higher Weil–Petersson volumes of moduli spaces of stable n-pointed curves. Comm. Math. Phys. **181** (3): 763–787.

[33] Kazarian, M., Lando, S. (2007). An algebro-geometric proof of Witten's conjecture. Journal of the American Mathematical Society, **20** (4): 1079–1089.

[34] Kontsevich, M. (1992). Intersection theory on the moduli space of curves and the matrix Airy function. Comm. Math. Phys., **147** (1): 1–23.

[35] Lavoie, J.-L., Grondin, F., Rathie, A. K.; Arora, K. (1994). Generalizations of Dixon's theorem on the sum of a ${}_3F_2$. Math. Comp., **62** (205): 267–276.

[36] Liu, K., Xu, H. (2009). A proof of the Faber intersection number conjecture. Journal of Differential Geometry, **83** (2): 313–335.

[37] Liu, K., Xu, H. (2011). The n-point functions for intersection numbers on moduli spaces of curves. Advances in Theoretical and Mathematical Physics, **15** (5): 1201–1236.

[38] Liu, K., Xu, H. (2008). Recursion formulae of higher Weil–Petersson volumes. International Mathematics Research Notices.

[39] Liu, S.-Q. (2015). From Witten conjecture to DVV's loop equation. unpublished.

[40] Macdonald, I. G. (1995). Symmetric functions and Hall polynomials (Vol. 354). Oxford: Clarendon press.

[41] Manin, Y. I., Zograf, P. (1999). Invertible cohomological field theories and Weil–Petersson volumes. arXiv preprint math/9902051.

[42] Mehta, M. L. (2004). Random matrices (Vol. 142). Academic press.

[43] Mirzakhani, M. (2007). Simple geodesics and Weil–Petersson volumes of moduli spaces of bordered Riemann surfaces. Inventiones mathematicae, **167** (1): 179–222.

[44] Mirzakhani, M. (2007). Weil–Petersson volumes and intersection theory on the moduli space of curves. Journal of the American Mathematical Society, **20** (1): 1–23.

[45] Mirzakhani, M. (2013). Growth of Weil–Petersson Volumes and Random Hyperbolic Surface of Large Genus. Journal of Differential Geometry, **94** (2): 267–300.

[46] Mulase, M., Safnuk, B. (2006). Mirzakhani's recursion relations, Virasoro constraints and the KdV hierarchy. Indian Journal of Mathematics **50** (2008): 189-228.

[47] Mumford, D. (1983). Towards an enumerative geometry of the moduli space of curves. In Arithmetic and geometry (pp. 271–328). Birkhäuser Boston.

[48] Okounkov, A. (2002). Generating functions for intersection numbers on moduli spaces of curves. International Mathematics Research Notices, (2002) **18**: 933–957.

[49] Okounkov, A., Pandharipande, R. (2009). Gromov-Witten theory, Hurwitz numbers, and matrix models. In I, Proc. Symposia Pure Math (Vol. 80, pp. 325–414).

[50] Sato, M. (1981). Soliton Equations as Dynamical Systems on a Infinite Dimensional Grassmann Manifolds (Random Systems and Dynamical Systems). RIMS Kokyuroku, **439**: 30–46.

[51] Segal, G., Wilson, G. (1985). Loop groups and equations of KdV type. Publications Mathmatiques de l'IHÉS, **61** (1): 5–65.

[52] Witten, E. (1991). Two-dimensional gravity and intersection theory on moduli space. Surveys in Diff. Geom., **1**: 243–310.

[53] Zhou, J. (2012). Intersection numbers on Deligne–Mumford moduli spaces and quantum Airy curve. arXiv preprint arXiv:1206.5896.

[54] Zhou, J. (2013). Topological Recursions of Eynard–Orantin Type for Intersection Numbers on Moduli Spaces of Curves. Letters in Mathematical Physics, **103** (11): 1191–1206.

[55] Zhou, J. (2013). Explicit formula for Witten–Kontsevich tau-function. arXiv preprint arXiv:1306.5429.

[56] Zhou, J. (2015) On Absolute N-Point Function Associated with Gelfand-Dickey polynomials. preprint.

[57] Zograf, P. (2006). An algorithm for computing Weil–Petersson volumes of moduli spaces of curves. Institut Mittag–Leffler Preprints.

Marco Bertola

Department of Mathematics and Statistics, Concordia University, 1455 de Maisonneuve W., Montréal, Québec, H3G 1M8, Canada

SISSA, via Bonomea 265, Trieste 34136, Italy

Centre de recherches mathématiques, Université de Montréal, C. P. 6128, succ. centre ville, Montréal, Québec, H3C 3J7, Canada

marco.bertola@concordia.ca, mbertola@sissa.it, bertola@crm.umontreal.ca

Boris Dubrovin

SISSA, via Bonomea 265, Trieste 34136, Italy

dubrovin@sissa.it

Di Yang

SISSA, via Bonomea 265, Trieste 34136, Italy

dyang@sissa.it